



March and August 2010 Population Size Estimates of *Oncorhynchus mykiss* in the Lower Tuolumne River

Prepared for
Turlock Irrigation District
333 East Canal Drive
Turlock, CA 95380

and

Modesto Irrigation District
1231 11th St
Modesto, CA 95354

Prepared by
Stillwater Sciences
2855 Telegraph Ave., Suite 400
Berkeley, CA 94705

November 2010

Stillwater Sciences. 2010. March and August 2010 population size estimates of *Oncorhynchus mykiss* in the Lower Tuolumne River. Prepared for the Turlock Irrigation District and the Modesto Irrigation District by Stillwater Sciences, Berkeley, CA. November.

SUMMARY

In both early-March and mid-August 2010, population size estimates of *Oncorhynchus mykiss* were developed in the lower Tuolumne River in accordance with the 3 April 2008 Delegated Order issued by the Federal Energy Regulatory Commission (FERC) implementing elements of a study plan previously developed in coordination with California Dept. of Fish and Game (CDFG), National Marine Fisheries Service (NMFS) and U.S. Fish and Wildlife Service (USFWS) biologists, and submitted to FERC on 16 July 2007.

Snorkel surveys were conducted during daylight hours from 1 to 8 March and from 17 to 24 August 2010 to estimate *O. mykiss* population size within the Tuolumne River. In addition to snorkel survey observations of *O. mykiss*, data for Chinook salmon (*O. tshawytscha*) and other species was also collected. Snorkel surveys were conducted using a two-phase survey design to sample five different habitat strata (i.e., riffle, run head, run body/tail, pool head, and pool body/tail) found downstream of La Grange Dam at river mile (RM) 51.8 using habitat typing from surveys performed in June 2008 (ending at RM 39.5) and March 2009 (from RM 39.5 down to RM 29.0). The study reaches extended from RM 51.8 to RM 38.4 near a bridge crossing within the 7-11 gravel operation in March and August 2010. A total of 66 of 181 sampling units in the study reach upstream of RM 38.4 were selected for either single pass or multi-pass snorkel surveys in July 2010. A total of 61 sampling units from the same study reach were selected for either single pass or multi-pass snorkel surveys in March 2010.

O. mykiss population estimates

Based upon the maximum count obtained over all dive passes in each sampled unit, only one young-of-the-year (YOY)/juvenile (< 150 mm FL) and 13 adult (> 150 mm FL) (sum total of 14) *O. mykiss* were observed in March 2010. During the August 2010 surveys, 313 YOY/juvenile (<150 mm FL) and 324 adult (> 150 mm FL) (sum total of 687) *O. mykiss* were observed along the study reach. Using a bounded counts population estimator (BCE) for the March 2010 survey period, a total of approximately 109 adult *O. mykiss* were present within the study reach (RM 51.8–38.4). No estimate was made for juvenile *O. mykiss* due to low count of only one individual. Using the same estimator for August 2010 survey period, approximately 2,405 juvenile and 2,139 adult *O. mykiss* were present within the study reach (RM 51.8–38.4).

The August 2010 juvenile *O. mykiss* population estimate of 2,405 was lower than the July 2009 estimate of 3,475 and similar to the July 2008 estimate of 2,472 juveniles. However, the summer population estimates are within the 95% CI for juvenile *O. mykiss* in all three years (2008-2010). The August 2010 adult *O. mykiss* population estimate of 2,139 was higher than both the July 2009 estimate of 963 and the July 2008 estimate of 643.

Chinook salmon population estimates

For Chinook salmon encountered during the March and August 2010 snorkel surveys, a maximum count of 577 juveniles (< 150 mm FL) were observed during March 2010 within all habitat types along the study reach and a maximum count of 1,028 juvenile Chinook salmon were observed in all habitat types during the August 2010 survey. This corresponded to bounded count population estimates of 6,141 Chinook salmon during the March 2010 surveys, and 6,338 during August 2010. By comparison, the July 2009 juvenile population estimate of 29,389 was much higher and the July 2008 estimate of 2,636 was lower. There were also 14 adult salmon observed in August 2010 as compared with 6 observed in July 2009, and 2 in July 2008.

Other species

A combination of native minnows (hardhead and Sacramento pikeminnow), along with native Sacramento sucker accounted for approximately 97% of non-salmonid fish observed for both the March and August sampling periods, with very low counts of non-native centrarchid species (largemouth bass, smallmouth bass) observed. Native minnows and suckers were found throughout the reaches in both sampling periods.

Relationship between Temperature and *O. mykiss* habitat use

To test the hypothesis that the summertime distribution of suitable habitat by observed life stages of *O. mykiss* is related to ambient river water temperature, water temperature data from thermographs deployed in the Tuolumne River were compared to juvenile and adult *O. mykiss* density from the August 2010 survey along the study reach. The data show that temperatures increased in the downstream direction, from 12.0°C (53.6°F) to 17.8°C (64.1°F) (maximum weekly average temperature [MWAT]), and that *O. mykiss* density of both adult and juveniles generally decreased along this same gradient. Although the longitudinal distribution of *O. mykiss* was similar for both the March and August surveys, the lower number of *O. mykiss* observations in March 2010, coupled with low water temperatures (maximum observed <14.5 °C [58.1 °F]) precluded any meaningful associations with temperature for the March 2010 surveys.

O. Mykiss habitat use at Restoration sites

A second hypothesis that habitat use by *O. mykiss* juveniles and adults observed in the Tuolumne River occurred at the same density in both restored and nearby reference sites was tested based on observed densities of *O. mykiss* juveniles and adults in habitat types (riffle, run head, and pool head) common to both groups in the August survey. For juveniles, this comparison showed riffle habitat use at upstream restoration sites was slightly greater than that of other riffle habitats. Juvenile habitat use within run head habitats was similar or reduced at the restoration sites in comparison to reference sites, with relatively low use of pool head habitat. For adults, this comparison showed a potential reduction of habitat use of riffle habitat at restoration sites, with similar use of run head habitat, and insufficient data for a comparison of pool head habitats.

Comparison with August 2010 Reference Survey Results

A comparison was made of *O. mykiss* and juvenile Chinook data collected during the August 2010 survey to “reference count”, snorkel survey data collected during August 2010 by TID/MID. The comparison shows a similar longitudinal trend, with overall densities decreasing in the downstream direction for both species. Along the study reach common to both surveys, a total of 195 *O. mykiss* juveniles and 73 adults were observed in the August reference count snorkel survey, while 210 juveniles and 253 adults were observed in the August BCE survey. A total of 142 juvenile Chinook were seen in the August reference survey with 889 seen in the August 2010 BCE survey.

Table of Contents

SUMMARY	II
1 INTRODUCTION	1
2 METHODS	2
2.1 Habitat Characterization	2
2.1.1 Habitat mapping	2
2.1.2 Habitat data collection.....	3
2.2 Snorkel Surveys	4
2.2.1 Study design and survey unit selection	4
2.2.2 Snorkel data collection	5
2.3 Water Quality and Flow	5
2.4 Water and Air Temperatures.....	6
2.5 Data analysis	6
2.5.1 Bounded counts population estimate.....	6
2.5.2 Comparisons with August 2010 TID/MID snorkel surveys.....	7
3 RESULTS	7
3.1 Habitat Characterization	7
3.1.1 March 2010	7
3.1.2 August 2010	8
3.2 Water Quality and Flow.....	8
3.2.1 March 2010	8
3.2.2 August 2010	9
3.3 Water and Air Temperature	10
3.3.1 March 2010	10
3.3.2 August 2010	11
3.4 Snorkel Surveys	12
3.4.1 March 2010	12
3.4.2 August 2010	19
4 DISCUSSION	26
4.1 Bounded Counts Study Assumptions.....	26
4.2 Variations in <i>O. mykiss</i> Population Estimates	27
4.2.1 March 2009	27
4.2.2 August 2010 and July 2008-9.....	27
4.3 <i>O. mykiss</i> Distribution in Relation to Water Temperature.....	28
4.3.1 March 2010	28
4.3.2 August 2010	28
4.4 Habitat Associations of <i>O. mykiss</i> and Chinook salmon Observations	29
4.4.1 March 2010	29
4.4.2 August 2010	31
4.5 Habitat Use at Restored and Reference Sites by <i>O. mykiss</i> and Chinook salmon	32
4.6 Comparison to August 2010 TID/MID Snorkel Surveys.....	33
4.6.1 <i>O. mykiss</i> observations	35
4.6.2 Chinook salmon observations	35
5 REFERENCES	36

List of Tables

Table 2-1. Coarse-scale habitat types used during snorkel surveys.	2
Table 2-2. Habitat data collected at each unit.	3
Table 2-3. Sample unit selection and survey count for March 2010.	4
Table 2-4. Sample unit selection and survey count for August 2010.	5
Table 2-5. Fish data collected within each unit during snorkel surveys.	5
Table 2-6. Water quality data collected during snorkel surveys.	6
Table 3-1. Summary of habitat types from RM 51.8 to 38.4, March and August 2010.	8
Table 3-2. Range of water quality data collected at snorkel sites during fish surveys in March 2010.	9
Table 3-3. Range of water quality data collected at snorkel sites during fish surveys in July 2009.	9
Table 3-4. Maximum weekly average temperature, seven-day average of daily maximum temperatures, and instantaneous maximum temperatures recorded by thermographs in the survey reach of the lower Tuolumne River during March 2010.	10
Table 3-5. Daily average, minimum, and maximum air temperature recorded at the NWS station at the Modesto Airport during the March 2010 snorkeling study period.	11
Table 3-6. Maximum weekly average temperature, seven-day average of daily maximum temperatures, and instantaneous maximum temperatures recorded by thermographs in the survey reach of the lower Tuolumne River during August 2010.	11
Table 3-7. Daily average, minimum, and maximum air temperature recorded at the NWS station at the Modesto Airport during the August 2010 snorkeling study period.	12
Table 3-8. Maximum count of <i>O. mykiss</i> by sampling unit, March 2010.	13
Table 3-9. Maximum count of <i>O. mykiss</i> by habitat type, March 2010.	14
Table 3-10. <i>O. mykiss</i> March 2010 bounded count population estimates between RM 51.8 and 38.4 by fish length and habitat type.	15
Table 3-11. Maximum counts of juvenile Chinook salmon by size class and sampling unit, March 2010.	16
Table 3-12. Maximum counts of juvenile Chinook salmon by size class and habitat type, March 2010.	17
Table 3-13. Chinook salmon March 2010 bounded count population estimates between RM 51.8 and 38.4 by fish length and habitat type.	18
Table 3-14. Maximum counts of non-salmonid species by sampling unit, March 2010.	18
Table 3-15. Maximum count of <i>O. mykiss</i> by NSO, August 2010.	20
Table 3-16. Maximum count of <i>O. mykiss</i> by habitat type, August 2010.	21
Table 3-17. <i>O. mykiss</i> August 2010 bounded count population estimates by fish length and habitat type.	22
Table 3-18. Maximum counts of juvenile Chinook salmon by size class and sampling unit, August 2010.	23
Table 3-19. Maximum counts of juvenile Chinook salmon by size class and habitat type, August 2010.	24
Table 3-20. Chinook salmon August 2010 bounded count population estimates by fish length and habitat type.	25
Table 3-21. Maximum counts of non-salmonid species by sampling unit (NSO), August 2010.	26
Table 4-1. Cover and substrate type found in sampling units with <i>O. mykiss</i> present during the March 2010 snorkel surveys.	30
Table 4-2. Cover and substrate type found in sampling units with Chinook salmon present during the March 2010 snorkel surveys.	30
Table 4-3. Cover and substrate type found in sampling units with <i>O. mykiss</i> present during the August 2010 snorkel surveys.	31

Table 4-4. Cover and substrate type found in sampling units with Chinook salmon present during the August 2010 snorkel surveys.....	31
Table 4-5. Salmonid observations in August reference and August BCE surveys in 2010 within the reach sampled during both studies.	34
Table 4-6. Salmonid counts and estimated densities in August reference and August BCE surveys in 2010 for units snorkeled during both dates.....	34

List of Figures

- Figure 1. Survey reach for March and July 2009 *O. mykiss* snorkel surveys in the lower Tuolumne River.
- Figure 2a. Hourly water and daily average air temperature and flow for the study reach from 1 February to 31 March 2010.
- Figure 2b. Hourly water and daily average air temperature and flow for the study reach from 1 July to 31 August 2010.
- Figure 3. Longitudinal distribution of major habitat type areas by river mile in the lower Tuolumne River (RM 52–40) for 2010 surveys.
- Figure 4a. Longitudinal distribution of major habitat type areas by river mile in the lower Tuolumne River (RM 52–40) for March 2010 surveys.
- Figure 4b. Longitudinal distribution of major habitat type areas by river mile in the lower Tuolumne River (RM 52–40) for August 2010 surveys.
- Figure 5. Size distribution of *O. mykiss* observed in Tuolumne River snorkel surveys, August 2010.
- Figure 6a. Distribution of observed *O. mykiss* counts among habitat types, by size class.
- Figure 6b. Distribution of observed *O. mykiss* density based on maximum count among habitat types, by size class.
- Figure 7. August 2010 adult *O. mykiss* density by river mile based upon maximum count in sampling units of each habitat type.
- Figure 8. August 2010 juvenile *O. mykiss* density by river mile based upon maximum count in sampling units of each habitat type.
- Figure 9. Longitudinal distribution of observed *O. mykiss* and water temperature in the lower Tuolumne River, August 2010.
- Figure 10. Observed densities of *O. mykiss* in individual sampling units in the August 2010 surveys.
- Figure 11. Observed densities of Chinook salmon in individual sampling units in the March 2010 surveys.
- Figure 12. Observed densities of Chinook salmon in individual sampling units in the August 2010 surveys.

List of Appendices

- Appendix A. Study Plan for 2009 surveys.
- Appendix B. 2009 Habitat maps.
- Appendix C. 2004 Habitat maps (McBain & Trush 2004).
- Appendix D. Habitat data.
- Appendix E. Water quality data.
- Appendix F. Water temperature data.
- Appendix G. Fish observation data.

1 INTRODUCTION

Routine fisheries monitoring surveys for the Don Pedro Project (FERC Project No. 2299) by the Turlock Irrigation District (TID) and Modesto Irrigation District (MID) have long documented the presence of *Oncorhynchus mykiss* in the lower Tuolumne River (TID/MID 2005). Summer snorkel surveys, conducted in most years since 1988, have documented an increased *O. mykiss* presence and relative abundance that is associated with the more consistent and higher summer flows provided since 1997 (TID/MID 2008).

On 19 March 1998, the National Marine Fisheries Service (NMFS) first listed the Central Valley steelhead as threatened under the Endangered Species Act (ESA). After several court challenges, NMFS issued a new final rule relisting the Central Valley steelhead on 5 January 2006 (71 FR 834). In a separate process resulting from terms of the 1996 FERC license amendment for the Project, NMFS staff provided input to a draft limiting factors analysis for Tuolumne River salmonids (Mesick et al. 2007) and included recommendations for developing abundance estimates, habitat use surveys, and anadromy determination of resident *O. mykiss*. These recommendations were conceptually used to develop the Districts' FERC Study Plan (TID/MID 2007), which was the subject of a 3 April 2008 FERC Order. As part of the Order, the Districts were required to conduct population estimate surveys in winter (February/March) and summer (June/July), with the first surveys starting in summer 2008 to determine *O. mykiss* population abundance by habitat type.

The Districts first submitted a detailed *O. mykiss* population estimate study plan (Stillwater Sciences 2008a) to FERC on 3 July 2008 to provide information on the abundance and habitat requirements within the lower Tuolumne River. A report on the July 2008 population size estimate (Stillwater Sciences 2008b) was submitted as part of the Districts' 2008 annual report to FERC (TID/MID 2009). An updated study plan (Stillwater Sciences 2009) was prepared in 2009 for the population estimate surveys and is attached to this report as Appendix A. In addition to providing data to develop population size estimates under current conditions, the study plan examined the following hypotheses:

Hypothesis 1: Summertime distribution of suitable habitat by observed life stages of *O. mykiss* is related to ambient river water temperature.

Hypothesis 2: Habitat use by *O. mykiss* juveniles and adults observed in the Tuolumne River occurs at the same density in both restored and nearby reference sites.

The *O. mykiss* snorkel surveys employed a two-phase sampling approach for the development of a reach-wide population estimate (Hankin and Mohr 2001) in the lower Tuolumne River. Survey sites were selected using a stratified random sampling approach, where the strata were major habitat types. In both March and August 2010, the overall sampling "universe" from which sampling strata were delineated extended from near La Grange Dam at river mile (RM) 51.8 to RM 38.4 at a bridge crossing within the 7-11 Materials, Inc. gravel operation (Figure 1). This reach coincides with the downstream areas where *O. mykiss* were observed (Riffle 31 at RM 38.0) during the August 2010 "reference count" snorkel surveys (Kirihara 2010).

The two-phase stratified sampling design involved snorkeling pre-selected sampling units (e.g., riffle, run, pool, etc.) multiple times in order to quantify the variance associated with density and subsequent population estimates. As in a typical Phase I sampling approach, primary snorkel surveys (Edmundson et al. 1968, Hankin and Reeves 1988, McCain 1992, Dolloff et al. 1996)

were conducted across a subset of the all sampling units. In Phase II, approximately 20–70% of each habitat type sampled was randomly selected for replicated surveys by repeated dive counts.

The methods presented by Stillwater Sciences (2009) discussed using a combined approach of both repeated dive counts and electrofishing. Current ESA permit restrictions for NMFS Section 10(a)(1)(A) permit No. 1282 (Stillwater) did not allow sufficient incidental take to conduct the second-phase surveys using electrofishing. Consequently, the surveys used only snorkel surveys, as provided for in the 2007 study plan and identified in letters provided by the Districts to FERC dated 3 July 2008 and 31 March 2009.

2 METHODS

2.1 Habitat Characterization

2.1.1 Habitat mapping

Habitat maps were compiled from an analysis of past habitat surveys, historical and more recent aerial photographs, and field surveys conducted in 2008, with results superimposed within a geographic information system (GIS). Field maps for the March and August 2010 BCE snorkel surveys were created using an orthorectified aerial photo and accompanying Light Detection and Ranging (LiDAR) topographic data from 21 September 2005 recorded at river flows of 321 cfs. Preliminary sampling unit boundaries of common habitat features (pools, riffles, and runs) were estimated from the LiDAR and bathymetric data between RM 52–38 within GIS by calculating locations corresponding to major water depth transitions (Table 2-1).

Table 2-1. Coarse-scale habitat types used during snorkel surveys.

Habitat type	Description ^a	Approximate depth
Riffle	Shallow with swift flowing, turbulent water. Partially exposed substrate dominated by cobble or boulder. Gradient moderate (less than 4%).	0–4 ft
Run	Fairly smooth water surface, low gradient, and few flow obstructions. Mean column velocity generally greater than one foot per second (fts ⁻¹).	4–10 ft
Pool	Slow flowing, tranquil water with mean column water velocity less than 1 fts ⁻¹ .	>10 ft

^a Major habitat types determined based upon observed hydraulic conditions (McCain 1992, Thomas and Bovee 1993, Cannon and Kennedy 2003)

As an initial validation of these coarse scale habitat types, we compared the habitat types mapped in July 2008 (Appendix B) with previous habitat type maps (Appendix C) developed by McBain and Trush (2004) between 1999–2001 on a base-layer map corresponding to a wetted perimeter of 622 cfs flown on 20 May 20 1991. Appendix C shows major habitat types (i.e., riffle, run, pool) encountered during the 1999–2001 surveys along with past and planned gravel introduction locations included in the *Tuolumne River Coarse Sediment Management Plan* (McBain and Trush 2004).

In general, habitat typing shown by McBain and Trush (Appendix C) indicates larger proportions of “pool” habitat types than those determined during this effort (Appendix B), which reserved the pool habitat designation for water depths greater than 10 ft. Additionally, because *O. mykiss* tend to congregate at transitions between habitat types, Appendix B shows a further division of pool and run body habitats into smaller, transitional habitat sampling units (pool head, pool tail, run head, and run tail) based upon location of slope channel slope break at the upstream and downstream end of the unit. For both the March and August 2010 surveys, pool tail and run tail habitats were consolidated into corresponding upstream pool body or run body habitat. This action was based on low use of the pool tail and run tail habitats as discrete sampling units in prior surveys (July 2008 and March 2009) and results in a reduced number of sampling units having low potential for use by salmonids available for habitat selection, thereby increasing the number of sampling units having a higher potential use, while not eliminating them from the area surveyed (see Section 2.2.1 for a complete description of sampling unit selection).

2.1.2 Habitat data collection

Float surveys were conducted in July 2008 and February 2009 to further refine and validate the preliminary habitat maps (Appendix B) described above at flows of approximately 106 cfs and 168 cfs, respectively. In addition to refining the locations and sizes of potential habitat sampling units, we collected habitat data (Table 2-2) at several locations within each sampling unit. Starting at upstream end of the study reach just downstream of La Grange Dam (Figure 1), habitat units were assigned a natural sequence order (NSO), a number, beginning with NSO 001, and incremented this identifier at each habitat transition (e.g., NSO 001 pool head, NSO 002 pool body, etc). The upstream and downstream end of each unit was located and marked on field maps, the location recorded with a handheld GPS unit, and labeled with flagging indicating the date, unit number, and habitat type.

Table 2-2. Habitat data collected at each unit.

Parameter	Method	Metric/Descriptor	Method reporting limit
Natural Sequence Order (NSO – Habitat unit #)	N/A	NSO-1, NSO-2, NSO-3, ...	N/A
Latitude/Longitude	Handheld GPS receiver	UTM	N/A
Habitat type	Visual estimation	See Table 2-1	N/A
Average unit width	Horizontal distance	Meters (feet) (measured at multiple transects)	0.01 m (0.1 ft)
Average unit length	Horizontal distance	Meters (feet)	0.01 m (0.1 ft)
Maximum/minimum depth	Vertical distance	Meters (feet)	0.15 m (0.5 ft)
Bed substrate composition	Visual estimation	Bedrock, boulder, cobble, gravel, organic, sand, silt	10%
Cover type	Visual estimation	None, boulder, cobble, IWM, bedrock ledges, overhead vegetation, aquatic vegetation	10%

Note that although the base layer of the 2009 habitat maps corresponds to a 2005 air photo at flows of 321 cfs, in order to provide a more accurate channel edge boundary for the March and July 2009 surveys, the channel edge of the habitat unit boundaries shown in Appendix B correspond to a wetted perimeter of 230 cfs previously digitized from air photos taken in 1986-87 and later refined to adjust for channel migration. The average daily flow during the March 2010 sampling was 224 cfs, and the average daily flow during the August 2010 sampling was 293 cfs. Because the estimated wetted perimeter of the habitat unit boundaries did not vary more than a few feet in most cases at these two flows, the channel edge boundary for 230 cfs was used for both the March and August 2010 surveys. For each habitat unit shown, habitat unit length and width were subsequently determined in GIS. Appendix D shows accompanying field habitat data collected in all habitat units mapped, including maximum depth and average width (usually at 1/3 and 2/3 of the unit's length), bed substrate composition, and instream cover type.

2.2 Snorkel Surveys

2.2.1 Study design and survey unit selection

After habitat typing and collecting habitat data in all units, a subset of units of each habitat strata was selected for single-pass snorkel surveys. The survey units were selected to balance the habitat sampling unit replication, total available number of units to draw from, coverage of at least 10% of the total length of a given habitat type, as well as sampling effort. The selection process involved random selection of one of the most upstream units of each habitat type, followed by a systematic uniform sampling of the remaining units in the study reach. After the first dive pass was completed, a tab was then pulled to determine if the unit was included in the second phase of sampling.

For the March 2010 surveys, a subset of 6–7 units was selected for each of the 5 habitat types, with the exception of the riffle habitat type for which 10 units were selected to capture habitat use at particular gravel augmentation projects (Table 2-3). In August 2010, a subset of 6–7 sampling units was selected from each of 5 habitat types (Table 2-4), with representative riffle habitats corresponding to restoration sites at some locations.

Table 2-3. Sample unit selection and survey count for March 2010.

Habitat	Phase I dives		Phase II survey	
	Initial units	Passes	Repeat units	Passes
Riffle	10	1	3	2
Pool head	6	1	3	2
Pool body /tail	6	1	3	2
Run head	7	1	3	2
Run body /tail	7	1	3	2
Total	36		30	

Table 2-4. Sample unit selection and survey count for August 2010.

Habitat	Phase I dives		Phase II survey	
	Initial units	Passes	Repeat units	Passes
Riffle	7	1	3	2
Pool head	6	1	3	2
Pool body /tail	6	1	3	2
Run head	6	1	3	2
Run body /tail	6	1	3	2
Total	31		30	

2.2.2 Snorkel data collection

Snorkel surveys were conducted during daylight hours from 1 to 8 March and 17 to 24 August 2010, respectively. A two-phase survey design was used to survey the various riffle, run, and pool strata. For the first phase, single-pass dive surveys were conducted by a four-person team. Sampling units were sampled from downstream to upstream in dive lanes using a zigzag pattern, passing fish and allowing them to escape downstream of the diver. If fish were observed to escape upstream, the diver took care to avoid counting these individuals twice. Divers recorded the type, length, and number of fish (Table 2-5). Total lengths were estimated in 50 mm size ranges (called “bins”) using markings on dive slates to correct for underwater size distortion.

Table 2-5. Fish data collected within each unit during snorkel surveys.

Parameter	Method	Metric/Descriptor	Method reporting limit
Date; start and end time	N/A	Day/month/year; hour/minute	N/A
Number of individuals	Visual estimation	Number	1
Fish length	Visual estimation	Millimeter	50 mm bins

The second phase of sampling required the collection of repeat dive counts and fish size data during each of two subsequent passes through the selected habitat units. These data were later used to statistically expand the dive counts to total population estimates for each habitat type. The Phase 2 dive pass replication was established at 2 passes in 2009 surveys to reduce sampling effort within particular sampling units while increasing the overall sample unit coverage (Stillwater 2010). Lastly, the occurrence of other non-salmonid native and non-native fish species was recorded as presence/absence and abundance.

2.3 Water Quality and Flow

At fish sampling locations, in addition to noting the type, length, and number of fish (Section 2.2), we collected spot measurements of *in situ* water quality data (temperature, dissolved oxygen, and conductivity) using a pre-calibrated multi-probe (YSI 85, Yellow Springs Instruments, Yellow Springs, OH) (Table 2-6). Dissolved oxygen (DO) probes were recalibrated each day and checked for accuracy in the laboratory against DO concentrations measured in

aerated tap water. Changes in underwater visibility were monitored horizontally using a Secchi disk oriented both toward and away from the sun. Daily average flow data for each day were obtained from the stream gage below the La Grange powerhouse at RM 51.8 (USGS No. 11289650).

Table 2-6. Water quality data collected during snorkel surveys.

Parameter	Method	Metric/Descriptor	Method reporting limit
Temperature	EPA 170.1	°C	0.1 °C
Dissolved oxygen	SM 4500-O	mg/L	0.01 mg/L
Conductivity	SM 2510A	umhos/cm	1.0 umhos/cm
Visibility	Secchi depth	meters (feet)	0.01 m (0.1 ft)

2.4 Water and Air Temperatures

From Spring 1987 to present, TID/MID has collected water temperature data from various locations in the lower Tuolumne River using recording thermographs (Hobo Pro V2 thermographs, OnSet Computer Corporation, Bourne, MA). The thermographs measured and stored water temperature data at one-hour intervals, with data downloads occurring at least twice a year.

Water temperature data collection during March and August 2010 also included spot measurements taken during snorkel surveys. The measurements were recorded over the course of the day as divers moved further downstream; as such, it was anticipated that these water temperatures would not be as representative as hourly thermograph recordings. The data do provide a general description of relative temperature conditions during dive surveys, however.

Regional air temperature data were obtained from the National Weather Service (NWS) station at Modesto Airport near RM 18. Water and air temperature data for the February through March, and July through August 2010 periods are presented in this report (Figures 2a and 2b).

2.5 Data analysis

2.5.1 Bounded counts population estimate

Water quality and fish observation counts were summarized by habitat unit type with initial density estimates calculated based upon the area searched within each habitat unit sampled. In addition to comparisons of fish density between habitat types, the density estimates and uncertainties were propagated across the unsampled areas for an overall reach-wide population estimate.

Population estimates were made for each stratum and size class using the general methods of Hankin and Mohr (2001). For units receiving multiple dives, the bounded counts formulae are used to produce an estimate of the unit population and an estimate of the variance of this estimate. Specifically, when there are r passes, and the counts of these are sorted in increasing order as $m_1 \leq m_2 \leq \dots \leq m_r$, the population is estimated as

$$\tilde{y}_B = m_r + (m_r - m_{r-1}),$$

and the mean squared error of this is estimated as

$$\text{M}\tilde{\text{S}}\text{E}(\tilde{y}_B) = (m_r - m_{r-1})^2.$$

The total population of multiply dived units is estimated as the sum of the bounded-counts estimates for the individual units. The total population of the survey region is estimated by expanding this, first to *all* dived units (singly or multiply dived) on the basis of mean dive counts, and then to all units (dived or undived) on the basis of area. An estimator of the variance of this is constructed from estimates of the mean-squared errors of the bounded-counts estimates for the multiply dived individual units, and the variance of the bounded-counts estimates around their common mean. The final formulae are included in Hankin and Mohr (2001). A nominal confidence interval for each stratum and size class was calculated formally as

$\hat{Y} \pm 1.96\sqrt{\hat{V}}$, where \hat{Y} and \hat{V} are the mean and variance estimates, *except* that the lower bound of this interval was “trimmed” to the number of fish actually observed.

2.5.2 Comparisons with August 2010 Reference Count snorkel surveys

Data collected during the August 2010 snorkel surveys (17–24 August) were compared to reference count snorkel survey data collected during 10–12 August 2010 (Kiriwara 2010). Although the sampled areas of these surveys differ, these data were collected only a few weeks prior to the data collected for this report, allowing for a general comparison of presence/absence and the relative proportions of larger and smaller size classes of *O. mykiss* and Chinook salmon in sampling units sampled during both surveys. Further, although TID/MID has sampled the same locations since 2001, we limit our comparison to the August 2010 data as these are the most directly comparable. There were no reference count survey data available for comparison with the March 2010 snorkel surveys.

3 RESULTS

3.1 Habitat Characterization

3.1.1 March 2010

For the total reach surveyed in March 2010 (RM 51.8–38.4), “run body/tail” habitat type occupied the greatest length of channel along the study reach, followed by riffles (Table 3-1). The “pool body/tail” habitat type, while less abundant than other habitat types (e.g., run head), occupied the third greatest length of channel. Other transitional habitat types (e.g., run head and pool head) accounted for only 4.8 % of the total reach length. Habitat maps and data for the entire study reach are shown in Appendices B and D. The longitudinal distribution of the area of each of the major habitat types within bins of 2 river miles is shown in Figure 3. Figure 4a presents the distribution of each of the major habitat types sampled in March 2010.

Table 3-1. Summary of habitat types from RM 51.8 to 38.4, March and August 2010.

Habitat type	Count	% by count	Total length (ft)	Total length (mi)	% reach length	Area (ft ²)
Riffle	40	22.1	15,271	2.89	21.4	1,281,867
Pool head	7	3.9	712	0.13	1.0	61,958
Pool body/tail	11/7	9.9	9,238	1.75	12.9	1,143,736
Run head	38	21.0	2,712	0.51	3.8	253,658
Run body/tail	42/36	43.1	43,423	8.22	60.9	4,449,862
Total	181	100.0	71,356	13.51	100.0	7,191,081

3.1.2 August 2010

The total reach surveyed in August 2010 (RM 51.8–34.8), was identical to the reach surveyed in March 2010 and therefore contains the same overall distribution of habitat types as shown in Table 3-1. Habitat maps and data for the entire study reach are shown in Appendices B and D. The longitudinal distribution of the area of each of the major habitat types within equal segments of 2 river miles is shown in Figure 3. Figure 4b presents the distribution of each of the major habitat types sampled in August 2010.

3.2 Water Quality and Flow

As water quality data were collected exclusively within units chosen for snorkel survey, data are presented by river mile, rather than by sampling unit, or summarized for the entire reach (Table 3-2 and Table 3-3). Water quality data for sampling units selected for snorkel surveys are shown in Appendix E.

Because of the strong influence of ambient air temperatures (Sullivan et al. 1990), temperatures of water released from the cold water pool of Don Pedro Reservoir increase in a downstream direction for both the spot measurements (Table 3-3) and in the continuous thermograph record during both the March and July survey periods (Appendix F). Note that the water temperature ranges shown in Table 3-2 and Table 3-3 represent changes over the course of the sampling day, and do not include nighttime temperatures or lows that are shown at representative thermograph locations in Appendix F.

3.2.1 March 2010

Daily average flow during the March 2010 survey period was 223 cfs. In general, dissolved oxygen concentration was high due to the low water temperatures. Horizontal visibility was reduced at the most downstream location due to local turbidity sources.

Table 3-2. Range of water quality data collected at snorkel sites during fish surveys in March 2010.

River miles	Sample date	Flow (cfs)¹	Water temp °C [°F]	DO (mg/L)	Horizontal visibility (ft)	Specific conductivity (uS/cm)
51.6–50.8	1 March	224	10.6–11.3 [51.1–52.3]	10.6–12.4	13.5	29.1–30.5
50.6–49.7	2 March	223	10.6–11.0 [51.1–51.8]	10.6–11.5	17	28.1–32.5
49.6–48.0	3 March	224	10.2–10.6 [50.4–51.1]	9.9–11.2	15	29.3–31.1
45.9	5 March	224	10.6 [51.1]	10.4	10.5	37.4
45.0–43.0	6 March	223	10.7–12.3 [51.3–54.1]	10.6–11.9	8.5–12	37.4–40.6
42.9–38.9	7 March	224	11.5–14.1 [52.7–57.4]	10.8–12.3	9–11.5	39.9–53.4
38.8–38.5	8 March	224	12.1–12.4 [53.8–54.3]	10.7–11.1	8.5	48.9–49.1

¹ Daily average flow data are measured from the stream gauge below La Grange powerhouse at RM 51.8 (USGS No. 11289650).

3.2.2 August 2010

Daily average flow during the August 2010 survey period ranged from 287–295 cfs. In general, there were only relatively small variations in water quality parameters at this flow range. Horizontal and vertical visibility indicated very low turbidity during the survey period.

Table 3-3. Range of water quality data collected at snorkel sites during fish surveys in July 2009.

River miles	Sample date	Flow (cfs)¹	Water temp °C [°F]	DO (mg/L)	Horizontal visibility (ft)	Specific conductivity (uS/cm)
51.8–51.6	17 August	293	12.6–12.6 [54.7–54.7]	9.8–9.8	32–32	30.4–30.4
50.8–50.3	18 August	287	12.7–13.1 [54.9–55.6]	11.0–11.2	27.3–31.5	28.8–29.1
49.9–49.7	19 August	294	14.3–14.3 [57.7–57.7]	11.3–11.3	27.3–27.3	29.3–29.3
49.1–48.0	20 August	295	14.2–16.4 [57.6–61.5]	11.2–13.1	25–25	29.4–29.7
46.9–45.1	21 August	294	13.9–15.3 [57.0–59.5]	11.8–12.7	20.5–20.5	30.4–31.1
45.0–43.2	22 August	293	13.3–15.4 [55.9–59.7]	10.9–11.2	19.0–21.5	31.5–32.0
42.7–39.6	23 August	293	15.6–18.5 [60.1–65.3]	11.3–12.0	16.5–15.5	33.2–37.1
39.2–38.8	24 August	293	16.3–16.3 [61.3–61.3]	9.7–9.7	17.5–17.5	38.2–38.2

¹ Daily average flow data are measured from the stream gauge below La Grange powerhouse at RM 51.8 (USGS No. 11289650).

3.3 Water and Air Temperature

The daily average water temperature for all thermographs and the daily minimum, maximum, and average air temperature (from the NWS station at the Modesto Airport) are shown in Appendix F. The range of daily averages, instantaneous maximum temperature, maximum weekly average temperature (MWAT), and the seven-day average of daily maximum temperature (7dayMAX) for the 1–8 March and 17–24 August study periods was determined, and all three metrics for both periods showed a similar trend of increasing in the downstream direction. The MWAT is the seven-day rolling average of average daily temperatures, and describes ambient water temperature conditions over the previous week. It is a standard used in water quality studies and total maximum daily load (TMDL) estimations of allowable temperature. The 7dayMAX is the seven-day rolling average of the daily maximum temperatures, and is a potentially more accurate indicator of conditions affecting survival and growth of salmonids (Sullivan et al. 2000, Stillwater Sciences 2002).

3.3.1 March 2010

During the March 2010 survey period, water temperature data collected by thermographs followed similar trends to spot temperature data collected during snorkel surveys, showing an increase in the downstream direction (Table 3-4). Along the study reach, the MWAT increased from 10.6°C (51.1°F) at Riffle A7 to 12.1°C (53.7°F) at the Ruddy Gravel site (Table 3-4). The 7dayMAX temperature ranged from 11.1°C (52.0°F) at the Riffle A7 location to 13.2°C (55.7°F) at the Ruddy Gravel site. The hourly, mean weekly average (MWAT), and 7dayMAX water temperatures for Riffle A7 (RM 50.8), Riffle 13B (RM 45.5), Roberts Ferry Bridge (RM 39.6), and Ruddy Gravel (RM 36.5) from 1 February to 31 March 2010 are presented graphically in Appendix F.

Table 3-4. Maximum weekly average temperature, seven-day average of daily maximum temperatures, and instantaneous maximum temperatures recorded by thermographs in the survey reach of the lower Tuolumne River during March 2010.

Monitoring location	RM	MWAT °C [°F] (week ending)	7dayMAX °C [°F] (week ending)	Instantaneous maximum °C [°F] (date)
Riffle A7	50.8	10.6 [51.1] (2 March)	11.1 [52.0] (2 March)	11.4 [52.5] (1 March)
Riffle 13B	45.5	11.3 [52.3] (7 March)	12.2 [54.0] (6 March)	12.7 [54.8] (1 March)
Roberts Ferry Bridge ¹	39.6	11.8 [53.3] (7 March)	12.8 [55.0] (7 March)	13.7 [56.7] (7 March)
Ruddy Gravel	36.5	12.1 [53.7] (7 March)	13.2 [55.7] (7 March)	14.2 [57.5] (7 March)

Note: Thermographs used have a reported error of ±0.2°C.

¹ Thermograph located approximately 0.75 miles upstream of bridge.

The average daily Modesto Airport air temperatures over the study period ranged from 8.3 to 12.8 °C (47.0 to 55.0 °F) with a high temperature of 18.9 °C (66.0 °F) (Table 3-5). The warmest day of March occurred after the study period on 17 March with an average daily temperature of 17.8 °C (64.0 °F) (Figure 2a) and a daily high temperature of 23.9 °C (75.0 °F). The highest daily maximum temperature in March occurred on 28 March with a reading of 26.1 °C (79.0 °F).

Table 3-5. Daily average, minimum, and maximum air temperature recorded at the NWS station at the Modesto Airport during the March 2010 snorkeling study period.

Date	Average air temperature °C [°F]	Minimum air temperature °C [°F]	Maximum air temperature °C [°F]
1 March 2009	12.8 [55]	6.7 [44]	18.3 [65]
2 March 2009	11.7 [53]	7.8 [46]	15.0 [59]
3 March 2009	8.9 [48]	6.7 [44]	11.1 [52]
4 March 2009	10.0 [50]	5.6 [42]	14.4 [58]
5 March 2009	8.3 [47]	2.2 [36]	13.9 [57]
6 March 2009	12.8 [55]	8.3 [47]	17.2 [63]
7 March 2009	12.2 [54]	5.0 [41]	18.9 [66]
8 March 2009	10.0 [50]	5.6 [42]	14.4 [58]

Hourly water temperature for several monitoring stations along the length of the study reach and daily air temperature from the Modesto Airport station was compared (Figure 2a). With flow being stable throughout period, Figure 2a shows that at the upstream-most monitoring station, water and air temperature are more independent of each other than at thermographs located farther downstream. That is, water temperature becomes more influenced by air temperature in the downstream direction, with water and air temperature peaks and troughs occurring at the same times of day at the downstream monitoring site at Roberts Ferry Bridge (RM 39.6).

3.3.2 August 2010

During the August 2010 survey period, water temperature data collected by thermographs followed similar trends to spot temperature data collected during snorkel surveys, which showed a general increase in the downstream direction (Table 3-6). Along the study reach, the MWAT increased from 12.0 °C (53.6 °F) at Riffle A7 to 17.8°C (64.1 °F) at Ruddy Gravel (Table 3-6). The 7dayMAX temperature ranged from 13.3°C (55.9 °F) at the Riffle A7 location to 19.2°C (66.6 °F) at the Roberts Ferry Bridge. The hourly, mean weekly average (MWAT), and 7dayMAX water temperatures for Riffle A7 (RM 50.8), Riffle 13B (RM 45.5), Roberts Ferry Bridge (RM 39.6), and Ruddy Gravel (RM 36.5) from 1 July to 31 August 2010 are presented graphically in Appendix F.

Table 3-6. Maximum weekly average temperature, seven-day average of daily maximum temperatures, and instantaneous maximum temperatures recorded by thermographs in the survey reach of the lower Tuolumne River during August 2010.

Monitoring location	RM	MWAT °C [°F] (week ending)	7dayMAX °C [°F] (week ending)	Instantaneous maximum °C [°F] (date)
Riffle A7	50.8	12.0 [53.6] (19 August)	13.3 [55.9] (19 August)	13.4 [56.0] (17 August)
Riffle 13B	45.5	14.5 [58.1] (18 August)	16.5 [61.7] (18 August)	16.7 [62.0] (17 August)
Roberts Ferry Bridge ¹	39.6	17.1 [62.7] (19 August)	18.5 [65.3] (18 August)	18.7 [65.6] (17 August)
Ruddy Gravel	36.5	17.8 [64.1] (19 August)	19.2 [66.6] (19 August)	19.5 [67.0] (17 August)

Note: Thermographs used have a reported error of ±0.2°C.

¹ Thermograph located approximately 0.75 miles upstream of bridge.

The average daily Modesto Airport air temperatures over the study period ranged from 21.7 to 27.8 °C (71.0 to 82.0 °F) with a high temperature of 38.9 °C (102 °F) (Table 3-7). The warmest day of August occurred just after the study period on 25 August with an average daily temperature of 30.0 °C (86 °F) and a daily high temperature of 41.7 °C (107 °F) (Figure 2b).

Table 3-7. Daily average, minimum, and maximum air temperature recorded at the NWS station at the Modesto Airport during the August 2010 snorkeling study period.

Date	Average air temperature °C [°F]	Minimum air temperature °C [°F]	Maximum air temperature °C [°F]
17 August 2010	24.4 [76.0]	16.1 [61.0]	32.8 [91.0]
18 August 2010	22.8 [73.0]	14.4 [58.0]	31.1 [88.0]
19 August 2010	24.4 [76.0]	14.4 [58.0]	33.9 [93.0]
20 August 2010	25.6 [78.0]	16.1 [61.0]	34.4 [94.0]
21 August 2010	21.7 [71.0]	14.4 [58.0]	28.9 [84.0]
22 August 2010	21.7 [71.0]	12.8 [55.0]	30.0 [86.0]
23 August 2010	24.4 [76.0]	14.4 [58.0]	34.4 [94.0]
24 August 2010	27.8 [82.0]	16.1 [61.0]	38.9 [102.0]

Hourly water temperature for several monitoring stations along the length of the study reach and daily air temperature from the Modesto Airport station was compared (Figure 2b). High flows through July kept water temperatures relatively low with little variability. Flow reductions in early August to approximately 300 cfs, shows a slight increase in variability among the temperature stations, but a continuation of relatively low temperatures, with a reduced influence of air temperature at thermographs located farther downstream (Figure 2b).

3.4 Snorkel Surveys

3.4.1 March 2010

3.4.1.1 *O. mykiss* observations

During the March 2010 survey period, divers observed 15 *O. mykiss* ranging from 0–600 mm (50 mm size bins) based upon maximum counts of all dive passes in each sampling unit (Table 3-8, Table 3-9 and Appendix G). These included one fish classified as a juvenile in the 50–99 mm size category, with the other 14 observed in the adult (>150 mm) size classes (Table 3-8 and Table 3-9). The *O. mykiss* were observed in 9 different sampling units from RM 51.6 to RM 38.5. The *O. mykiss* were observed in all habitat types, with the exception of the “Run body/tail” habitat, with the juvenile observation in a pool head habitat unit at RM 51.6 (Table 3-8 and Table 3-9).

Table 3-8. Maximum count of *O. mykiss* by sampling unit, March 2010 (data are divided into 50 mm total length size classes).

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	0-49 mm	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm	>500 mm
51.6	4	Pool head	Y	1								1		
51.6	5/6	Pool body/tail	Y									2		2
50.9	11	Pool body	N											
50.8	12/13	Run body/tail	N											
50.6	15	Run head	Y								1			
50.5	16/17	Run body/tail	Y											
50.3	18	Riffle	N											
50.3	19	Run head	N										1	
50.1	20/21	Run body/tail	N											
50.1	22	Riffle	Y											
49.7	26	Riffle	Y						2					
49.7	27	Pool head	N											
49.6	28/29	Pool body/tail	Y									1		
48.8	42	Run head	Y											
48.7	43/44	Run body/tail	N											
48.0	54	Pool head	N											
45.9	70	Riffle	N											
45.0	86	Pool head	Y											
44.8	90	Run head	N											
44.7	93	Riffle	Y											
44.5	101	Riffle	N											
43.7	104	Pool body	N											
43.0	111	Riffle	N											
43.0	112	Pool head	Y							2				
43.0	113/114	Pool body/tail	Y											
42.9	116/117	Run body/tail	Y											
42.9	119	Run head	N											

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	0-49 mm	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm	>500 mm
42.3	126	Riffle	N								1			
41.9	133	Run head	Y											
41.8	134/135	Run body/tail	N											
39.2	165	Pool head	N											
38.9	166/167	Pool body/tail	N											
38.9	168	Riffle	N											
38.8	172	Run head	N											
38.7	173/174	Run body/tail	Y											
38.5	179	Riffle	N									1		
Total (maximum unit count of all passes)				1	0	0	0	0	2	2	2	5	1	2

Table 3-9. Maximum count of *O. mykiss* by habitat type, March 2010 (data are divided into 50 mm total length size classes).

Habitat	0-49 mm	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm	>500 mm	Total (max. unit count of all passes)
Pool body/tail									3		2	5
Pool head	1						2		1			4
Riffle						2		1	1			4
Run body/tail												0
Run head								1		1		2
Totals by size class	1	0	0	0	0	2	2	2	5	1	2	15

3.4.1.2 *O. mykiss* population estimate

Table 3-10 shows the March 2010 *O. mykiss* population estimate for the lower Tuolumne River by length (<150 mm for YOY and juvenile; >150 mm for adults) and habitat type using the method of bounded counts (Hankin and Mohr 2001) for the study reach from RM 51.8 to RM 38.4 . Since the YOY/juvenile observations of *O. mykiss* were minimal (n=1), no population estimate for this lifestage was derived from the March 2010 survey. From an observed 13 adult *O. mykiss* in March 2010, an estimated population of 109 adults (with a 95% CI of 50-168) was determined (Table 3-10). Adult *O. mykiss* were observed in all habitat types with the exception of “run body/tail” habitat.

Table 3-10. *O. mykiss* March 2010 bounded count population estimates between RM 51.8 and 38.4 by fish length and habitat type.

Habitat	<i>O. mykiss</i> < 150 mm				<i>O. mykiss</i> ≥ 150 mm			
	Obs. ¹	Est.	St. dev.	95% CI ²	Obs.	Est.	St. dev.	95% CI ²
Pool head	1	1	0.3	1–2	3	6	2.6	3–11
Pool body/tail	0	--	--	--	4	14	6.2	4–26
Riffle	0	--	--	--	4	37	14.1	9–64
Run head	0	--	--	--	2	53	25.6	3–103
Run body/tail	0	--	--	--	0	--	--	--
Total	1	1	0.3	1–2	13	109	30.0	50–168

¹ Largest numbers seen in any single dive pass for each unit, summed over units. Note that because of the potential for the same fish to be assigned to different size classes on subsequent passes, summation of the largest numbers assigned to individual (50 mm) size bins yields may overestimate total fish observed.

² Nominal confidence intervals calculated as + 1.96 standard deviations.

3.4.1.3 Chinook salmon observations

Table 3-11 and Table 3-12 show the number of Chinook salmon observed within the study reach during the March 2010 surveys, based on the maximum count by pass, resulting in a total of 577 observations. All Chinook salmon were YOY and juveniles found within the 0–49 and 50–99 mm size classes. These salmon were seen in 16 different sampling units ranging from RM 51.6 to RM 38.8 (Table 3-11) and all habitat types (Table 3-12).

Table 3-11. Maximum counts of juvenile Chinook salmon by size class and sampling unit, March 2010.

River mile	Sampling unit	Habitat type	Multiple pass survey (Y/N)	0–49 mm	50–99 mm
51.6	4	Pool head	Y	18	
51.6	5/6	Pool body/tail	Y	76	
50.9	11	Pool body	N		
50.8	12/13	Run body/tail	N		
50.6	15	Run head	Y		
50.5	16/17	Run body/tail	Y		
50.3	18	Riffle	N	172	9
50.3	19	Run head	N		
50.1	20/21	Run body/tail	N	80	
50.1	22	Riffle	Y	8	
49.7	26	Riffle	Y		1
49.7	27	Pool head	N		
49.6	28/29	Pool body/tail	Y		
48.8	42	Run head	Y		
48.7	43/44	Run body/tail	N		
48.0	54	Pool head	N		
45.9	70	Riffle	N	41	25
45.0	86	Pool head	Y		
44.8	90	Run head	N		
44.7	93	Riffle	Y	6	16
44.5	101	Riffle	N	1	
43.7	104	Pool body	N		
43.0	111	Riffle	N	2	
43.0	112	Pool head	Y	15	15
43.0	113/114	Pool body/tail	Y		
42.9	116/117	Run body/tail	Y	23	44
42.9	119	Run head	N		
42.3	126	Riffle	N	2	10
41.9	133	Run head	Y		
41.8	134/135	Run body/tail	N	1	
39.2	165	Pool head	N		
38.9	166/167	Pool body/tail	N		
38.9	168	Riffle	N		
38.8	172	Run head	N	8	3

River mile	Sampling unit	Habitat type	Multiple pass survey (Y/N)	0–49 mm	50–99 mm
38.7	173/174	Run body/tail	Y	1	
38.5	179	Riffle	N		
Total (max. unit count of all passes)				454	123

Table 3-12. Maximum counts of juvenile Chinook salmon by size class and habitat type, March 2010.

Habitat	0–49 mm	50–99 mm	Total (maximum unit count of all passes)
Pool body/tail	76		76
Pool head	33	15	48
Riffle	232	61	293
Run body/tail	105	44	149
Run head	8	3	11
Totals by size class	454	123	577

No adult Chinook salmon were observed within the study reach. The complete Chinook salmon observation data by pass are shown in Appendix G.

3.4.1.4 Chinook salmon population estimate

Table 3-13 shows the March 2010 Chinook salmon population estimate for the lower Tuolumne River by length (<150 mm for YOY and juvenile; >150 mm for adults) and habitat type using the method of bounded counts (Hankin and Mohr 2001). Since there were no observations of adult Chinook salmon, no population estimate for this lifestage was derived from the March 2010 survey. From an observed 574 YOY/juvenile Chinook salmon in March 2010, an estimated population of 6,141 (with a 95% CI of 2,687–9,596) was determined (Table 3-10). Juvenile Chinook salmon were observed in all habitat types, with riffle habitat providing the highest number of observations and generating the largest portion of the population estimate (approx. 55%).

Table 3-13. Chinook salmon March 2010 bounded count population estimates between RM 51.8 and 38.4 by fish length and habitat type.

Habitat	Chinook salmon < 150 mm				Chinook salmon ≥ 150 mm			
	Obs. ¹	Est. ²	St. dev.	95% CI ³	Obs. ¹	Est. ²	St. dev.	95% CI ³
Pool head	48	67	22.2	48–111	0	--	--	--
Pool body/tail	76	238	153.8	76–540	0	--	--	--
Riffle	293	3,386	898.0	1,626–5,146	0	--	--	--
Run head	11	--	--	--	0	--	--	--
Run body/tail	146	2,449	1,508.7	146–5,406	0	--	--	--
Total	574	6,141	1,762.6	2,687–9,596	0	--	--	--

¹ Largest numbers seen in any single dive pass for each unit, summed over units. Note that because of the potential for the same fish to be assigned to different size classes on subsequent passes, summation of the largest numbers assigned to individual (50 mm) size bins yields may overestimate total fish observed.

² Estimate for run head habitat type for juvenile salmon not included in overall population estimate due to lack of multiple pass data to develop an expansion factor.

³ Nominal confidence intervals calculated as + 1.96 standard deviations.

3.4.1.5 Non-salmonid observations

Several other fish species were observed and counted during the March 2010 survey period (Table 3-14). Most other fish seen within the study reach were native species in the minnow (*Cyprinidae*) and sucker (*Catostomidae*) families. A combination of hardhead and Sacramento pikeminnow, along with Sacramento sucker accounted for 97.0%. Other observed non-salmonid fish included catfish (*Ictaluridae*), centrarchids (largemouth bass, smallmouth bass), and sculpin (*Cottidae*), accounted for the remaining 3% of observations. Most centrarchids occurred toward the downstream end of the study reach where water temperatures were slightly warmer, while native suckers were found throughout the reach. The complete non-salmonid fish observation data are in Appendix G.

Table 3-14. Maximum counts of non-salmonid species by sampling unit, March 2010.

RM	Sampling unit	Habitat	CF	LMB	SMB	SC	HH/PM	SS
50.8	12/13	Run body/tail						1
50.6	15	Run head						3
50.5	16/17	Run body/tail						35
50.3	18	Riffle						10
50.1	20/21	Run body/tail						10
50.1	22	Riffle						1
49.7	26	Riffle						4
49.7	27	Pool head						1
49.6	28/29	Pool body/tail				1		8
48.8	42	Run head						6
48.7	43/44	Run body/tail						8
48.0	54	Pool head					10	2
45.9	70	Riffle	1					4
44.7	93	Riffle					7	
44.5	101	Riffle						3
43.0	112	Pool head						3

RM	Sampling unit	Habitat	CF	LMB	SMB	SC	HH/PM	SS
43.0	113/114	Pool body/tail		1				
42.9	116/117	Run body/tail					2	3
42.3	126	Riffle						3
41.9	133	Run head						4
41.8	134/135	Run body/tail						19
38.9	166/167	Pool body/tail						1
38.7	173/174	Run body/tail			1	1		1
38.5	179	Riffle						10
Total (all sampled units)			1	1	1	2	19	140

CF = catfish species; LMB = largemouth bass; SMB = smallmouth bass; SC = sculpin species; HH/PM = hardhead/Sacramento pikeminnow; SS = Sacramento sucker

3.4.2 August 2010

3.4.2.1 *O. mykiss* observations

During the August 2010 survey period, divers observed 682 *O. mykiss* ranging from 0–500 mm (50 mm size bins) based upon maximum counts of all dive passes in each sampling unit (Table 3-15, Table 3-16). Approximately half of these fish (320) were YOY/juvenile (<150 mm), with a total of 362 adults (>150 mm) observed (Figure 5). Complete fish observation data by sampling unit and dive pass is presented in Appendix G.

The *O. mykiss* were observed in 22 different sampling units from RM 51.8 to RM 39.7 and in all habitat types (Table 3-15 and Table 3-16). Habitat use and reach-wide distribution of YOY/juvenile and adult *O. mykiss* were similar, based on the maximum count from dive passes (Figure 6a) highest in riffle and run body/tail habitats. Fish densities (Figure 6b) for juvenile size classes (<150 mm) highest in riffle and pool head habitats. Juvenile size classes were also observed in each of the other habitat types, with lowest density in pool body habitats (Figure 6b). Adult-size classes (>150 mm) were observed in highest density in pool head habitats, with lower densities found in each of the other habitat types (Figure 6b).

Adult fish habitat use was concentrated at upstream sampling units (above RM 45.0) and primarily occurred at transitional run head and pool head habitats (Figure 7). Juvenile fish habitat use showed a similar distribution from upstream to downstream and occurred primarily at riffle habitat types, along with transitional run head and pool head habitat types (Figure 8).

Table 3-15. Maximum count of *O. mykiss* by sampling unit, August 2010 (data are divided into 50 mm total length size classes).

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm
51.8	1	Pool Head	Y		1		7	10	6	2	1	
51.6	4	Pool Head	Y					4	3	2	2	1
51.6	5	Pool body/tail	Y		2	2	5	2	4	1	2	1
50.8	12	Run body/tail	Y	50	23	13	2	12	24	10	1	
50.6	14	Riffle	Y	6	60	28	10	4	3	2		
50.3	19	Run Head	Y		6	5	5	3	7			
49.9	24	Run body/tail	N		7	4	1	2	13	4		
49.7	27	Pool Head	Y	3	7	12	2	1	1			
49.6	28	Pool body/tail	Y		2	4	2	8	5	3		
49.1	38	Run Head	N		1							
48.4	45	Riffle	N	9	26	5						
48.1	51	Run body/tail	Y		16	4	1	1	1	1		
48.0	53	Riffle	N		4					1		
48.0	54	Pool Head	N		6	5	1		3			
46.9	62	Run Head	Y		5	8	3		2	1		
45.3	81	Pool body/tail	N									
45.1	83	Run body/tail	N		13	9	3		5			
45.0	86	Pool Head	N		8	11	3	5	2			
44.8	90	Run Head	N									
44.5	101	Riffle	Y		15	13	2	1				
43.7	104	Pool body/tail	N									
43.2	107	Riffle	Y		19	8	3	1	2			
42.7	123	Run Head	N									
42.4	124	Run body/tail	Y	7	21	5		1				
40.3	150	Run body/tail	N		2	3	1					
39.7	156	Riffle	N		1	1						
39.6	157	Run Head	Y									
39.2	165	Pool Head	N									

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm
38.9	166	Pool body/tail	N									
38.9	168	Riffle	N									
38.8	171	Pool body/tail	Y									
Total (maximum unit count of all passes)				75	245	140	51	55	81	27	6	2

Table 3-16. Maximum count of *O. mykiss* by habitat type, August 2010 (data are divided into 50 mm total length size classes).

Habitat	50-99 mm	100-149 mm	150-199 mm	200-249 mm	250-299 mm	300-349 mm	350-399 mm	400-449 mm	450-499 mm	Total (max. unit count of all passes)
Pool body/tail		4	6	7	10	9	4	2	1	43
Pool head	3	22	28	13	20	15	4	3	1	109
Riffle	15	125	55	15	6	5	3			224
Run body/tail	57	82	38	8	16	43	15	1		260
Run head		12	13	8	3	9	1			46
Totals by size class	75	245	140	51	55	81	27	6	2	682

3.4.2.2 *O. mykiss* population estimate

Table 3-17 shows the August 2010 *O. mykiss* population estimate for the lower Tuolumne River by length (<150 mm for YOY and juvenile; >150 mm for adults) and habitat type using the method of bounded counts (Hankin and Mohr 2001). Out of an estimated 2,405 juveniles and 2,139 adults *O. mykiss* in August 2010 (an overall population estimate of 4,544), we estimated a 95% confidence interval of 625–4,185 and 727–3,552 for YOY/juvenile and adults, respectively (Table 3-17).

The relative differences between population estimates and observed fish counts are due to differences in habitat unit areas (e.g., run body/tail habitat types occupying approximately 20 times more habitat area than run head units (Table 3-2). This results in higher population estimates in some habitat types even though the observed counts may be similar or lower than those found in other habitat types. In August 2010, juvenile and adult population estimates were shown to be highest in run body/tail and riffle habitat types (Table 3-17).

Table 3-17. *O. mykiss* August 2010 bounded count population estimates by fish length and habitat type.

Habitat	<i>O. mykiss</i> < 150 mm				<i>O. mykiss</i> ≥ 150 mm			
	Obs. ¹	Est. ²	St. dev.	95% CI ³	Obs. ¹	Est.	St. dev.	95% CI ³
Pool head	24	42	8.4	26–58	72	90	6.3	78–102
Pool body/tail	4	12	4.9	4–22	32	136	109.5	32–351
Riffle	139	756	178.0	407–1,105	78	412	118.9	179–645
Run head	12	163	86.8	12–333	26	286	185.3	26–649
Run body/tail	134	1,432	886.2	134–3,169	116	1,215	677.3	116–2,542
Total	313	2,405	908.1	625–4,185	324	2,139	720.6	727–3,552

¹ Largest numbers seen in any single dive pass for each unit, summed over units. Note that because of the potential for the same fish to be assigned to different size classes on subsequent passes, summation of the largest numbers seen assigned to individual (50 mm) size bins may overestimate total fish observed.

² Estimate for *O. mykiss* juveniles in pool head habitats not included in overall population estimate due to lack of multiple pass data to develop an expansion factor.

³ Nominal confidence intervals calculated as + 1.96 standard deviations. Standard deviation and confidence intervals undefined for multiple pass units with identical dive counts.

3.4.2.3 Chinook salmon observations

Divers observed a large number of juvenile Chinook salmon within the study reach during August 2010 as well as small numbers within the adult size classes (>150 mm). Salmon were seen in 19 different sampling units from RM 51.8 to RM 31.9 (Table 3-18) and all habitat types (Table 3-19). Most salmon were juveniles found within the 50–99 mm size class.

Table 3-18. Maximum counts of juvenile Chinook salmon by size class and sampling unit, August 2010.

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	0-49 mm	50-99 mm	100-149 mm	150-199 mm	600-649 mm	650-699 mm	700-799 mm	900-999 mm
51.8	1	Pool head	Y						2	3	1
51.6	4	Pool head	Y								
51.6	5	Pool body/tail	Y		87						
50.8	12	Run body/tail	Y	148	29	14					
50.6	14	Riffle	Y	110	31	4					
50.3	19	Run head	Y	9	40	20		1			
49.9	24	Run body/tail	N	50	37	32	1				
49.7	27	Pool head	Y		3	1					
49.6	28	Pool body/tail	Y		3	1	4				
49.1	38	Run head	N								
48.4	45	Riffle	N	30	104	52					
48.1	51	Run body/tail	Y	14	22	4	2				
48.0	53	Riffle	N		4						
48.0	54	Pool head	N		2						
46.9	62	Run head	Y	10	27	10					
45.3	81	Pool body/tail	N								
45.1	83	Run body/tail	N		20	8					
45.0	86	Pool head	N								
44.8	90	Run head	N		1						
44.5	101	Riffle	Y	5	31	11					
43.2	107	Riffle	Y		18	3					
42.7	123	Run head	N								
42.4	124	Run body/tail	Y		19	11					
40.3	150	Run body/tail	N								
39.7	156	Riffle	N								
39.6	157	Run head	Y								
39.2	165	Pool head	N								
38.9	166	Pool body/tail	N			1					

RM	Sampling Unit	Habitat	Multiple pass survey (Y/N)	0-49 mm	50-99 mm	100-149 mm	150-199 mm	600-649 mm	650-699 mm	700-799 mm	900-999 mm
38.9	168	Riffle	N			2					
38.8	171	Pool body/tail	Y								
Total (maximum unit count of all passes)				376	478	174	7	1	2	3	1

Table 3-19. Maximum counts of juvenile Chinook salmon by size class and habitat type, August 2010.

Habitat	0-49 mm	50-99 mm	100-149 mm	150-199 mm	600-649 mm	650-699 mm	700-799 mm	900-999 mm	Total (max. unit count of all passes)
Pool body/tail		90	2	4					96
Pool head		5	1			2	3	1	12
Riffle	145	188	72						405
Run body/tail	212	127	69	3					411
Run head	19	68	30		1				118
Totals by size class	376	478	174	7	1	2	3	1	1,042

Divers observed a total of seven adult Chinook salmon (>600 mm) at two separate sampling units in the upper portion of the study reach at RM 51.8 and RM 50.3. A total of seven salmon in the 150–199 mm size class (a size class technically included as “adult”, but not typically observed) were seen in three separate sampling units between RM 49.9 and 48.1. The complete Chinook salmon observation data by pass are shown in Appendix G.

3.4.2.4 Chinook salmon population estimate

Table 3-20 shows the August 2010 Chinook salmon population estimate for the lower Tuolumne River by length (<150 mm for YOY and juvenile; >150 mm for adults) and habitat type using the method of bounded counts (Hankin and Mohr 2001). Out of an estimated 6,338 juveniles and 117 adult Chinook salmon in August 2010 (an overall population estimate of 6,455), we estimated a 95% confidence interval of 3,291–9,385 and 14–249 for YOY/juvenile and adults, respectively (Table 3-20). The data show that the greatest estimated abundance of YOY and juvenile Chinook salmon occurred in run body/tail and riffle habitats, with the greatest estimated abundance of adults in the run body/tail habitat type (Table 3-20).

Table 3-20. Chinook salmon August 2010 bounded count population estimates by fish length and habitat type.

Habitat	Chinook salmon < 150 mm				Chinook salmon ≥ 150 mm			
	Obs. ¹	Est.	St. dev.	95% CI ²	Obs. ¹	Est. ²	St. dev.	95% CI ³
Pool head	5	13	5.3	5–23	6	7	4.0	6–15
Pool body/tail	92	324	115.8	97–551	4	24	31.1	4–85
Riffle	400	2,149	571.2	1,029–3,268	0	--	--	--
Run head	97	1,054	606.0	97–2,242	1	20	25.4	1–70
Run body/tail	379	2,798	1,307.6	379–5,361	3	65	53.8	3–170
Total	973	6,338	1,554.6	3,291–9,385	14	117	67.3	14–249

¹ Largest numbers seen in any single dive pass for each unit, summed over units. Note that because of the potential for the same fish to be assigned to different size classes on subsequent passes, summation of the largest numbers assigned to individual (50 mm) size bins may overestimate total fish observed.

² Estimate adult salmon within riffle habitats for adult salmon not included in overall population estimate due to lack of multiple pass data to develop an expansion factor.

³ Nominal confidence intervals calculated as ± 1.96 standard deviations.

3.4.2.5 Non-salmonid observations

Several other fish species were observed during the August 2010 study period (Table 3-21). Most fish seen within the study reach were native species in the minnow (*Cyprinidae*) and sucker (*Catostomidae*) families. A combination of cyprinids (hardhead and Sacramento pikeminnow), along with Sacramento sucker accounted for 89.5% of observed non-salmonid fish. Non-native striped bass were observed in six sampling units (primarily pool body habitat) from RM 51.8 to RM 38.9. The complete non-salmonid fish observation data are in Appendix G.

Table 3-21. Maximum counts of non-salmonid species by sampling unit, August 2010.

RM	Sampling unit	Habitat	GAM	LP	LMB	HH/PM	SB	SCP	SMB	SS
51.8	1	Pool head				1	1			
51.6	5	Pool body/tail					2			
50.8	12	Run body/tail				5		3		64
50.6	14	Riffle						7		6
50.3	19	Run head		1			1			70
49.9	24	Run body/tail	100			40	1			100
49.7	27	Pool head								1
49.6	28	Pool body/tail								10
49.1	38	Run head	3							40
48.4	45	Riffle	3					2		
48.1	51	Run body/tail				8				24
48.0	53	Riffle						1		3
48.0	54	Pool head								3
46.9	62	Run head								4
45.3	81	Pool body/tail				7				24
45.1	83	Run body/tail				3				77
45.0	86	Pool head				15				1
44.8	90	Run head				1				
44.5	101	Riffle				31		1		14
43.7	104	Pool body/tail			1	1	7			180
43.2	107	Riffle				6				8
42.4	124	Run body/tail				41			1	147
40.3	150	Run body/tail				19				
39.7	156	Riffle				3				150
39.6	157	Run head				2				40
38.9	166	Pool body/tail			1	15	1		1	9
38.9	168	Riffle				1				
38.8	171	Pool body/tail								1
Total (all sampled units)			106	1	2	199	13	14	2	1313

GAM = Gambusia sp.; LP= Lamprey sp.; LMB = large mouth bass; HH/PM = heardhead/pikeminnow; SB = Striped bass; SCP = Sculpin sp.; SMB = small mouth bass; SS = Sacramento sucker

4 DISCUSSION

4.1 Bounded Counts Study Assumptions

It should be noted that the bounded counts method was developed for use in smaller stream systems (Hankin and Mohr 2001) and applying the methodology to a larger system such as the Tuolumne River is only feasible provided key assumptions are satisfied. One critical assumption of the bounded counts approach is that all individuals have an equal probability of being observed. As noted above, this assumption may be challenged in locations with large numbers of juvenile Chinook salmon, due to low visibility conditions in deeper pool habitats, as well as low visibility due to light and background turbidity variations within the river between seasons or

from upstream to downstream. For these reasons, the resulting population estimates may be low-biased.

A second assumption of the bounded counts method is that observation efficiency is not 100%, so the number of fish seen in any single dive pass is, in general, an underestimate of the true number of fish present. For a closed population where fish do not migrate into or out of the unit between dives, the maximum number of fish seen over multiple passes is a low-biased estimator of the true population. However, because larger habitat units were subsampled at some locations, for run habitat types in particular, the resulting density expansions may have introduced a high-biased estimate of the true population size since fish are able to migrate freely into and out of the searched area due to the lack of habitat boundaries relevant to the sampled fish (e.g., riffle transitions) in many locations.

4.2 Variations in *O. mykiss* Population Estimates

4.2.1 March Survey Period

Overall, the March 2010 population estimate of 109 adult *O. mykiss* (>150 mm) was low, with virtually no representation of juvenile size classes (<150 mm) relative to adults (Table 3-10). Although the high numbers of Chinook salmon juveniles observed during the March 2010 surveys (Table 3-12) may have resulted in misidentification of some *O. mykiss* within the same area, the low numbers of juvenile *O. mykiss* observed is consistent with a winter-spring spawning period that begins in February (Moyle 2002). The low number of adult *O. mykiss* observed are consistent with the results of the March 2009 survey. The low numbers of *O. mykiss* during spring were attributed to one or more of the following potential causes:

1. Adult *O. mykiss* have a heterogeneous (i.e., “patchy”) distribution and it appears that even though the 2010 winter sampling efforts were conducted in the same reach as summer surveys, upstream of Roberts Ferry Bridge (RM 39.5), the resulting observation of adults remains low. Information from other sources (e.g., from angling or tracking) may identify whether habitat use is distributed farther downstream.
2. Adult *O. mykiss* may be more furtive in winter, swimming into or occupying deeper portions of pools or out of range of the diver visibility, which is also reduced in winter due to lower light levels and increased turbidity. Nighttime dive surveys could be considered in future surveys, since low light situations tend to reduce the startle reflex of *O. mykiss*.
3. Lastly, adult *O. mykiss* may be altogether absent from the survey reach because they have migrated downstream of RM 29 or did not survive the previous over-summer conditions. This could be confirmed by any of: a) catch and release angling outside of the survey reach, b) capture, implantation of acoustic tags and tracking as provided in the TID/MID (2007) study plan, or c) video observations at the Districts Alaska type counting weir recently deployed at RM 24 in September 2009.

4.2.2 August Survey Period

The August 2010 population estimate of 4,544 *O. mykiss* indicates a relatively equal proportion of juveniles (2,405) relative to adults (2,139) (Table 3-17). In comparison to the July 2008 results of 2,472 juveniles and 643 adults, and the July 2009 results of 3,475 juveniles and 963 adults, the August 2010 results indicate a relatively similar number of juveniles over the 2008-2010 summer

sampling periods, and a noticeable increase in the number and proportion of adults. Juvenile *O. mykiss* population estimates would be expected to vary from year-to-year due to the large number of potential eggs deposited by each additional female spawner. Also, the juvenile estimates (Table 3-17) are all within the with 95% CIs computed from 2008-2010 (Stillwater Sciences 2008b, 2010).

The August 2010 adult *O. mykiss* population estimate may relate to conditions in the river below La Grange dam that were greatly influenced by flood control releases occurring from April thru July 2010. These releases extend cooler water temperatures farther downstream. In addition, flood bypass releases around the generating units at the Don Pedro powerhouse during May-June 2010 may have resulted in the introduction of *O. mykiss* into the river from upstream reservoirs. In August 2010, small groups of larger sized (>250 mm) adult *O. mykiss* were observed in run body and pool body habitats downstream of where they were observed in previous survey years (2008 and 2009). These adults appeared as similar in size, coloration, and condition and were observed schooling together in circular patterns. Larger numbers of smaller sized (150-200 mm) adult fish were also observed in August 2010 (Figure 5). These sized fish would not have been able to come from the 2010 year class and also indicate possible introduction from upstream reservoirs due to flood control releases.

4.3 *O. mykiss* Distribution in Relation to Water Temperature

4.3.1 March 2010

During the March 2010 snorkel surveys, water temperatures remained below 14.5°C throughout the study reach, with daily average temperatures exceeding 13.0°C only at the lowest sampling unit (RM 38.4) on 7 March 2010. These temperature conditions are not thought to particularly affect the distribution of *O. mykiss* and it is likely that some other factor may also explain the decreasing *O. mykiss* density with distance downstream of La Grange Dam. All *O. mykiss* observed were found at or upstream of RM 38.5, similar to the March 2009 survey. As discussed above in Section 4.2, presence/absence of *O. mykiss* downstream of the study reach could be confirmed by any of: a) catch and release angling outside of the survey reach, b) capture, implantation of acoustic tags and tracking as provided in the TID/MID (2007) study plan, or c) video observations at the Districts Alaska type counting weir deployed at RM 24 in September 2009. Counting weir results show only one adult *O. mykiss* (276 mm) detected during the operational period from September 22, 2009 through January 31, 2010 (TID/MID 2010). Preliminary results from an acoustic tag and tracking studying initiated by the Districts' in February 2010 are currently not available, pending completion of the study.

4.3.2 August 2010

To test Hypothesis #1 that summertime distribution of observed life stages of *O. mykiss* across suitable habitat is related to ambient river water temperature, we compared water temperature data taken from thermographs to fish density in the sampled units. The data show that temperatures increase in the downstream direction (Section 3.3.2, Table 3-6) and that the density of adult *O. mykiss* (>150 mm) generally decreased along this same gradient (Figure 9). In sampling units where fish were seen, density of adult fish was generally similar from just downstream of La Grange Dam to approximately RM 47, with a peak density near RM 45 (Figure 9). The density of adults then decreased markedly in the downstream direction. As noted in

Section 4.2.2, conditions in the river below La Grange dam were greatly influenced by flood control releases that extend cooler water temperatures farther downstream.

Similar to adults, the density of YOY and juvenile *O. mykiss* decrease in the downstream direction, with generally similar distribution from just downstream of La Grange Dam to approximately RM 43 (Figure 9). Peak density of juveniles occurred near RM 45, with very low densities below RM 43. Juveniles were found in six out of seven riffle sampling units, indicating a strong preference for this habitat type. However, juveniles were also observed in five out of six sampling units, although in lower density (Table 6a and Table 6b). Generally, juveniles were not expected in this habitat type at downstream locations for a number of reasons, including predation and territorial exclusion by the larger size classes of *O. mykiss*. The occurrence of juveniles in this habitat type may also have been related to the earlier flood control releases, where juveniles were simply displaced from an upstream habitat due to increased water velocity, or where physical habitat (e.g. depth, velocity, cover, food supply) became available as microhabitat along the stream margin of run habitats.

4.4 Habitat Associations of *O. mykiss* and Chinook salmon Observations

4.4.1 March 2010

Table 4-1 and Table 4-2 show the range of cover and substrate components observed during habitat mapping for each habitat type where *O. mykiss* and Chinook salmon were present during the March 2010 surveys. Variations in cover types and amounts were limited in all sampling units, with higher percentages of the “No Cover” class found throughout the reach (Appendix D-2). For this reason, the cover results do not provide a meaningful basis for establishing a relationship with habitat use by juveniles or adults of either species. Chinook salmon juveniles were the most observed salmonid during the surveys and were observed primarily in riffle and transitional pool head and run head habitats where higher percentages of cobble were reported (Table 4-1).

Table 4-1. Cover and substrate type found in sampling units with *O. mykiss* present during the March 2010 snorkel surveys.

Cover type	Pool body/tail	Pool head	Riffle	Run body/tail	Run head
Cover type range (%)					
Boulder	0–10	0–0	0–5	No fish observed	0–10
Wood	0–0	0–5	0–0		0–0
Ledge	0–0	0–0	0–0		0–0
Overhang	0–5	0–0	5–20		0–0
Aquatic vegetation	0–10	0–30	0–0		0–10
No cover	85–90	65–100	80–100		90–90
Substrate type range (% covering channel bed)					
Bedrock	20–50	0–50	0–0	No fish observed	0–0
Boulder	20–20	10–20	10–20		10–20
Cobble	25–40	30–50	50–60		50–60
Gravel	0–10	0–30	20–40		20–40
Sand	5–10	0–10	0–10		0–0
Silt	0–0	0–0	0–0		0–0
Organic	0–0	0–0	0–0		0–0

Table 4-2. Cover and substrate type found in sampling units with Chinook salmon present during the March 2010 snorkel surveys.

Cover type	Pool body/tail	Pool head	Riffle	Run body/tail	Run head
Cover type range (%)					
Boulder	0–0	0–0	0–5	0–0	0–0
Wood	0–0	0–5	0–0	0–0	0–0
Ledge	0–0	0–0	0–0	0–0	0–0
Overhang	0–0	0–0	5–20	0–5	0–5
Aquatic vegetation	0–10	0–30	0–5	0–0	0–0
No cover	90–90	65–100	80–95	95–100	95–95
Substrate type range (% covering channel bed)					
Bedrock	0–50	0–50	0–10	0–15	0–0
Boulder	0–20	10–20	10–20	10–20	0–0
Cobble	0–25	30–50	20–60	40–60	0–60
Gravel	0–0	0–30	20–70	20–30	0–30
Sand	0–5	0–10	0–10	0–10	0–10
Silt	0–0	0–0	0–0	0–0	0–0
Organic	0–0	0–0	0–0	0–0	0–0

4.4.2 August 2010

Table 4-3 and Table 4-4 show the range of cover and substrate components observed during habitat mapping for each habitat type where *O. mykiss* and Chinook salmon were present during the August 2010 surveys. As in March 2010, variations of cover types and amounts were limited in all sampling units, with higher percentages of sampling units with no cover found throughout the reach (Appendix D-2). Therefore cover results do not provide a meaningful basis for establishing a relationship with habitat use by juveniles or adults of either species. Nevertheless, *O. mykiss* and Chinook salmon were observed primarily in riffle and run body/tail habitats where higher percentages of cobble were reported relative to other substrates associated with those habitat types (Table 4-3).

Table 4-3. Cover and substrate type found in sampling units with *O. mykiss* present during the August 2010 snorkel surveys.

Cover type	Pool body/tail	Pool head	Riffle	Run body/tail	Run head
Cover type range (%)					
Boulder	0–10	5–10	0–10	0–5	0–0
Wood	0–0	0–5	0–5	0–5	0–5
Ledge	0–0	0–0	0–10	0–0	0–0
Overhang	0–5	5–10	5–10	5–10	5–10
Aquatic vegetation	0–10	0–0	0–5	0–50	0–10
No cover	85–90	85–100	80–95	35–100	90–90
Substrate type range (% covering channel bed)					
Bedrock	20–50	10–50	0–10	10–20	0–0
Boulder	0–20	10–50	10–20	10–60	10–20
Cobble	25–40	30–60	50–70	20–50	60–70
Gravel	0–10	5–30	20–40	10–40	0–20
Sand	5–10	5–10	0–10	10–20	0–0
Silt	0–0	0–0	0–0	0–0	0–0
Organic	0–0	0–0	0–0	0–0	0–0

Table 4-4. Cover and substrate type found in sampling units with Chinook salmon present during the August 2010 snorkel surveys.

Cover type	Pool body/tail	Pool head	Riffle	Run body/tail	Run head
Cover type range (%)					
Boulder	10–10	5–10	0–10	0–5	0–0
Wood	0–0	0–0	0–5	0–5	0–5
Ledge	0–0	0–0	0–10	0–0	0–0
Overhang	0–5	0–5	5–10	5–10	5–10
Aquatic vegetation	0–10	0–0	0–0	0–50	0–10
No cover	85–90	85–90	80–100	35–95	90–90

Cover type	Pool body/tail	Pool head	Riffle	Run body/tail	Run head
Substrate type range (% covering channel bed)					
Bedrock	20–50	10–20	10–10	0–10	0–0
Boulder	20–20	10–50	10–20	10–60	10–20
Cobble	20–40	40–60	50–70	20–50	40–70
Gravel	10–50	5–10	20–40	10–40	20–50
Sand	5–30	5–10	10–10	10–20	0–10
Silt	0–0	0–0	0–0	0–0	0–0
Organic	0–0	0–0	0–0	0–0	0–0

4.5 Habitat Use at Restored and Reference Sites by *O. mykiss* and Chinook salmon

Hypothesis #2 states that the density of *O. mykiss* juveniles and adults is the same in restored sites as in nearby reference sites in the Tuolumne River. This hypothesis was originally formulated with the intention of testing habitat use at planned gravel augmentation sites (TID/MID 2007). However, other than the CDFG gravel addition projects near Old La Grange Bridge, completed from 2001–2003, and the joint Tuolumne River Technical Advisory Committee/Friends of the Tuolumne (FOT) gravel augmentation at Bobcat Flat (RM 43) in 2005, no further gravel augmentation projects have been implemented since that time. This has limited the sampling replication and statistical power to detect any differences between restored and reference sites.

As a means to evaluate habitat use of these restoration sites, observed densities of *O. mykiss* juveniles and adults were compared at the three habitat types that were sampled within the restoration sites to the same habitat types surveyed elsewhere in August 2010. The low number of *O. mykiss* observations in March 2010 do not allow for meaningful comparisons. Figure 10 shows the *O. mykiss* density of juveniles and adults at pool head, riffle, and run head habitat types sampled in August 2010 from sampling units found at both the restoration sites and from all similar sample units within the study reaches upstream of RM 38.0. For juvenile *O. mykiss* the densities show a relatively high use of riffle habitat at restoration sites when compared with other riffle sampling units; with relatively similar use of run head habitat at the upstream restoration sites; and an overall low density in pool head habitats found at the downstream portion of the reach (Figure 10). These same patterns appear for adult *O. mykiss* the densities throughout the reach.

A similar evaluation was done using juvenile Chinook salmon. Figures 11 and 12 show juvenile Chinook densities as sampled in March 2010 and August 2010, respectively for the same three habitat types. In March 2010, juvenile Chinook densities at the restoration sites were greater in each of the habitat types when compared to the reference sampling units (Figure 11), with the exception of riffle habitats between RM 44–46. In August 2010, juvenile Chinook densities either exceeded or were similar to the reference units (Figure 12). Considering the similar habitat preferences for juvenile *O. mykiss* and juvenile Chinook salmon, it appears that salmonid use of restoration sites is similar, or possibly enhanced within riffle habitats, when compared with nearby reference sites. Additional replication through either an increased number of gravel augmentation sites, or an increased number of survey events would be needed to improve the statistical power enough to detect whether significant differences in habitat use exist.

4.6 Comparison to August 2010 Reference Count Snorkel Surveys

Results from the August 2010 snorkel data were compared to observations made during the August 2010 reference count snorkel survey (Kiriwara 2010) for the sampled reach common to both surveys and within sampling units surveyed during both sampling events (Table 4-5 and Table 4-6). The August 2010 BCE data are observations from the first pass of the multiple pass bounded count estimation method to allow a direct comparison to August 2010 reference survey, which came from single pass snorkel surveys that employ catch-per-unit-effort (CPUE) methodology. Note that the reference count surveys were not conducted in March, precluding comparison with the March 2010 surveys.

Table 4-5. Salmonid observations in August reference count (single pass) and August BCE (first pass) surveys in 2010 within the reach sampled during both studies.

August 2010 reference count snorkel survey					August 2010 BCE snorkel survey				
Location	RM	<150 mm <i>O. mykiss</i> count	>150 mm <i>O. mykiss</i> count	<150 mm <i>O.</i> <i>tshawytscha</i> count	Sampling Units	RM	<150 mm <i>O. mykiss</i> count	>150 mm <i>O. mykiss</i> count	<150 mm <i>O. tshawytscha</i> count
Riffle A7 – R31	50.7– 38.0	195	73	142	1–181	51.8–38.4	210	253	889

Table 4-6. Salmonid counts and estimated densities in August reference count (single pass) and August BCE (first pass) surveys in 2010 for units snorkeled during both dates.

Location	RM	August 2010 reference count snorkel survey									August 2010 BCE snorkel surveys								
		Site	Habitat type	Area (ft ²)	<150 mm <i>O. mykiss</i>		>150 mm <i>O. mykiss</i>		<150 mm <i>O.</i> <i>tshawytscha</i>		Samplin g Unit	Habitat type	Area (ft ²)	<150 mm <i>O. mykiss</i>		>150 mm <i>O. mykiss</i>		<150 mm <i>O.</i> <i>tshawytscha</i>	
					#	#/ft ²	#	#/ft ²	#	#/ft ²				#	#/ft ²	#	#/ft ²	#	#/ft ²
Riffle A7	50.6	1	Riffle	6,000	16	0.0133	0	0	20	0.186	14	Riffle	45,670	30	0.0007	34	0.0007	120	0.0026
Riffle 2	49.1	2	Pool- Run	6,000	13	0.0014	3	0.0014	16	0.019	28,29	Pool Body/ Tail	23,835	4	0.0002	9	0.0004	105	0.0044
Riffle 5B	46.9	3	Run- Pool	9,375	11	0.0012	1	0.0002	7	0.0007	54	Pool Head	14,569	2	0.0001	9	0.0006	1	0.0001

4.6.1 *O. mykiss* observations

A total of 195 *O. mykiss* juveniles and 73 adults were observed in August 2010 reference count survey, while 210 juveniles and 253 adults were observed in the August 2010 BCE survey (Table 4-5). The between-site comparison shows similar longitudinal trends for juveniles, with observations and density generally decreasing in the downstream direction (Table 4-6). In both surveys, the greatest abundance of *O. mykiss* juveniles occurred within riffle habitat near RM 50.6 (Table 4-6). Adult *O. mykiss* abundance was lower for the August reference survey when compared with the August BCE survey at shared sampling sites. This was particularly evident at the upstream riffle location near RM 50.6 where no adults were observed during the reference survey and 34 adults were observed during the BCE survey (Table 4-6).

It should be noted that the August 2010 reference count survey data were collected from sites established in past years and targeted based on prior years' data as likely areas of relatively high *O. mykiss* abundance. The area surveyed during the August BCE surveys was greater (by an order of magnitude in most cases) than in June (Table 4-6). The reference count snorkel survey reoccupies the same sampling units and areas on an annual basis, produces a yearly index with which to evaluate yearly trends, assuming reoccupied sampling units and areas are representative of the entire reach. The BCE methodology (Hankin and Mohr 2001) produces a population estimate, with appropriate confidence intervals, that, due to the incorporation of multiple passes in each unit and greater area searched in each unit and along the reach, can be used to evaluate habitat- and reach-wide distribution patterns.

4.6.2 Chinook salmon observations

A total of 142 Chinook salmon juveniles were observed during the August 2010 reference survey, while a total of 889 juveniles were observed during the August BCE survey (Table 4-5). As noted above, the total area in the BCE surveys is greater than in the reference surveys. Salmon were observed in each habitat type sampled by the two methods. Although a stream-type life history strategy is not believed to be common for Chinook salmon in the Tuolumne River, the presence of juveniles in mid-summer indicates that conditions (e.g., water temperature, food availability) in summer 2010 were suitable for survival in upper portions of the reach.

5 REFERENCES

- Cannon, T.C., and T. Kennedy. 2003. Snorkel survey of the lower American River 2003. Prepared by Fishery Foundation of California, San Francisco for U.S. Fish and Wildlife Service, Central Valley Project Improvement Program, Sacramento, California.
- Dolloff, A., J. Kershner, and R. Thurow. 1996. Underwater observations. Pages 533–551 *in* B. R. Murphy and D. W. Willis, editor. Fisheries techniques. American Fisheries Society, Bethesda, Maryland.
- Edmundson, E., F. E. Everest, and D. W. Chapman. 1968. Permanence of station in juvenile Chinook salmon and steelhead trout. *Journal of the Fisheries Research Board of Canada* 25: 1453–1464.
- Hankin, D.G. and M. Mohr. 2001. Improved two-phase survey designs for estimation of fish abundance in small streams. Preprint from David G. Hankin, Department of Fisheries Biology, Humboldt State University, Arcata, California.
- Hankin, D. G., and G. H. Reeves. 1988. Estimating total fish abundance and total habitat area in small streams based on visual estimation methods. *Canadian Journal of Fisheries and Aquatic Sciences* 45: 834–844.
- Kirihara, S. 2010. 2010 Lower Tuolumne River annual report. Report 2010-5: 2010 snorkel report and summary update. Prepared by Turlock and Modesto Irrigation Districts, California and Stillwater Sciences, Berkeley, California for Federal Energy Regulatory Commission, Washington, D.C., *in prep*.
- McCain, M. E. 1992. Comparison of habitat use and availability for juvenile fall Chinook salmon in a tributary of the Smith River, California. *FHR Currents*. No. 7. USDA Forest Service, Region 5.
- McBain and Trush. 2004. Habitat maps for the upper 15.8 miles of the gravel bedded reach. Appendix D *in* Coarse sediment management plan for the lower Tuolumne River. Revised final report. Prepared by McBain and Trush, Arcata, California for Tuolumne River Technical Advisory Committee, Turlock and Modesto Irrigation Districts, USFWS Anadromous Fish Restoration Program, and CALFED Bay Delta Authority.
- Mesick, C., J. McClain, D. Marston, and T. Heyne. 2007. Draft limiting factor analyses & recommended studies for fall-run Chinook salmon and rainbow trout in the Tuolumne River, attachment 2 to USFWS comments on FERC study plan. http://elibrary.FERC.gov/idmws/file_list.asp?accession_num=20070314-0089 [Accessed 10 October 2008].
- Moyle, P. B. 2002. *Inland fishes of California*. Revised edition. University of California Press, Berkeley.
- Stillwater Sciences. 2002. Stream temperature indices, thresholds, and standards used to protect coho salmon habitat: a review. Unpublished white paper. Prepared by Stillwater Sciences, Arcata, California for Campbell Timberland Management, Fort Bragg, California.

Stillwater Sciences. 2008a. Population size estimates of resident *O. mykiss* in the lower Tuolumne River. Study plan. Prepared by Stillwater Sciences, Berkeley, California for Turlock Irrigation District and Modesto Irrigation District.

Stillwater Sciences. 2008b. July 2008 population size estimate of *Oncorhynchus mykiss* in the Lower Tuolumne River. Prepared for the Turlock Irrigation District and the Modesto Irrigation District by Stillwater Sciences, Berkeley, CA. October

Stillwater Sciences. 2009. Study Plan for Population Size Estimates of *O. mykiss* in the lower Tuolumne River. Prepared by Stillwater Sciences, Berkeley, California for Turlock Irrigation District and Modesto Irrigation District. January

Stillwater Sciences. 2010. March and July 2009 population size estimates of *Oncorhynchus mykiss* in the Lower Tuolumne River. Prepared for the Turlock Irrigation District and the Modesto Irrigation District by Stillwater Sciences, Berkeley, CA.

Sullivan, K., D. J. Martin, R. D. Cardwell, J. E. Toll, and S. Duke. 2000. An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria. Draft report. Prepared for Sustainable Ecosystems Institute, Portland, Oregon.

Sullivan, K. J. J. Tooley, K. Doughty, J. E. Caldwell, P. Knudsen. 1990. Evaluation of prediction models and characterization of stream temperature regimes in Washington. Timber/Fish/Wildlife Report Number TFW-WQ3-90-006. Washington Department of Natural Resources, Olympia, Washington.

Thomas, J. A., and K. D. Bovee. 1993. Application and testing of a procedure to evaluate transferability of habitat suitability criteria. *Regulated Rivers: Research and Management* 8: 285–294.

TID/MID (Turlock Irrigation District/Modesto Irrigation District). 2005. Ten year summary report of Turlock Irrigation District and Modesto Irrigation District pursuant to Article 58 of the license for the Don Pedro Project, No. 2299. 1 Volume.

TID/MID. 2007. Tuolumne River fisheries study plan - Don Pedro Hydroelectric Project (FERC No. 2299). Prepared by T. Ford, N. Hume, S. Wilcox, and R. Yoshiyama for Turlock Irrigation District and Modesto Irrigation District.
http://elibrary.FERC.gov/idmws/file_list.asp?accession_num=20070718-0082 [Accessed 10 October 2008].

TID/MID. 2008. Report 2007-7: 2007 Rainbow trout data summary report. *In* 2007 report of Turlock Irrigation District and Modesto Irrigation District pursuant to Article 39 of the license for the Don Pedro Project, No. 2299.

TID/MID. 2009. 2008 report of Turlock Irrigation District and Modesto Irrigation District pursuant to Article 39 of the license for the Don Pedro Project, No. 2299.

TID/MID. 2010. Report 2009-8: 2009 Counting weir report. *In* 2010 report of Turlock Irrigation District and Modesto Irrigation District pursuant to Article 39 of the license for the Don Pedro Project, No. 2299.




Figure 1. BCE study reach on the lower Tuolumne River, March and August 2010.




Figure 2a. Hourly water temperature, daily average air temperature, and daily average flow for the study reach from 1 February to 31 March 2010.




Figure 2b. Hourly water temperature, daily average air temperature, and daily average flow for the study reach from 1 July to 31 August 2010.




Figure 3. Longitudinal distribution of major habitat type areas by river mile in the lower Tuolumne River (RM 52-30) for March and August 2010 surveys.




Figure 4a. Longitudinal distribution of major habitat type areas sampled by river mile in the lower Tuolumne River (RM 52-38) for March 2010 survey.




Figure 4b. Longitudinal distribution of major habitat type areas sampled by river mile in the lower Tuolumne River (RM 52-38) for August 2010 survey.





Figure 5. Size distribution of *O. mykiss* observed in Tuolumne River snorkel surveys, August 2010. For units receiving multiple passes, the count is from the pass with the largest count for that size class.

small fish (<150 mm)



large fish (≥ 150 mm)





Figure 6a. Distribution of observed *O. mykiss* counts among habitat types, by size class in August 2010. For units receiving multiple passes, the count is from the pass with the largest count.

small fish (<150 mm)



large fish (≥ 150 mm)




Figure 6b. Distribution of observed *O. mykiss* density based on maximum count among habitat types, by size class in August 2010.




Figure 7. August 2010 adult *O. mykiss* density by river mile based upon maximum count in sampling units of each habitat type.




Figure 8. August 2010 juvenile *O. mykiss* density by river mile based upon maximum count in sampling units of each habitat type.




Figure 9. Longitudinal distribution of observed *O. mykiss* and water temperature in the lower Tuolumne River, August 2010. Solid diamonds are observed zeros, open diamonds are observed non-zero values.




Figure 10. Observed densities of *O. mykiss* in individual sampling units in the March 2010 surveys. Densities are maximum dive counts (in parenthesis) divided by the area sampled. Restoration sites are shown with broken lines (7-11 [RM 39.0], FOT [RM 43.0], CDFG 2001 [RM 50.3], CDFG 2003 [RM 50.6]). Non-restoration sites are shown with solid lines.




Figure 11. Observed densities of *O. tshawytscha* in individual sampling units in the March 2010 surveys. Densities are maximum dive counts (in parenthesis) divided by the area sampled. Restoration sites are shown with broken lines (7-11 [RM39.0], FOT [RM 43.0], CDFG 2001 [RM 50.3], CDFG 2003 [RM 50.6]). Non-restoration sites are shown with solid lines.




Figure 12. Observed densities of *O. tshawytscha* in individual sampling units in the August 2010 surveys. Densities are maximum dive counts (in parenthesis) divided by the area sampled. Restoration sites are shown with broken lines (7-11 [RM 39.0], FOT [RM 43.0], CDFG 2001 [RM 50.3], CDFG 2003 [RM 50.6]). Non-restoration sites are shown with solid lines.

Appendices

Appendix A: Study Plan (2009)



Study Plan for Population Size Estimates of *O. mykiss* in the lower Tuolumne River

Prepared for
Turlock Irrigation District
333 East Canal Drive
Turlock CA 95380

and

Modesto Irrigation District
1231 11th St
Modesto, CA 95354

Prepared by
Stillwater Sciences
2855 Telegraph Ave. Suite 400
Berkeley, CA 94705

January 2009


Stillwater Sciences

Table of Contents

1	BACKGROUND AND PURPOSE.....	3
2	FIELD SAMPLING AND DATA COLLECTION	3
2.1	Habitat Typing	4
2.2	Sample Site Selection.....	5
2.3	Sampling Period.....	7
2.4	Measurement Parameters and Sampling Methods	7
2.4.1	Snorkel Surveys	8
2.4.2	Electrofishing at Riverine Sites	9
2.4.3	Fish Handling Protocols	10
2.5	Hypothesis Testing.....	10
2.6	Field Work Notification	10
3	QUALITY ASSURANCE.....	11
3.1	Data Quality Objectives for Measurement Data	11
3.2	Training Requirements/Certification	12
3.3	Instrument/Equipment Testing, Inspection, and Maintenance Requirements	12
3.4	Instrument Calibration and Frequency	12
3.5	Reconciliation with Data Quality Objectives	12
3.6	Data Management	13
4	DATA ANALYSIS.....	13
5	REPORTING	13
6	PERMITTING REQUIREMENTS	13
7	REFERENCES.....	15

List of Tables

Table 1.	Coarse scale habitat types to be used during snorkel surveys	4
Table 2.	Estimated number of sampling units that will meet study design assumption of sampling at least 10% of the total length of a given habitat type.	6
Table 3.	Measurement parameters and methods for snorkel surveys	8
Table 4.	Preliminary sample unit selection and survey count.....	9
Table 5.	Field Work Notification.....	11
Table 6.	Data quality objectives for field parameters	11

Appendices

Appendix A	Lower Tuolumne River Habitat Mapping and Habitat Types from RM 52-40
Appendix B	Preliminary Habitat Mapping and Habitat Types in the lower Tuolumne River from RM 40-30

1 BACKGROUND AND PURPOSE

Fisheries monitoring for the Don Pedro Project (FERC Project No. 2299) by the Turlock Irrigation District (TID) and Modesto Irrigation District (MID) has long documented the presence of *Oncorhynchus mykiss* (*O. mykiss*) in the lower Tuolumne River (TID/MID 2005). On March 19, 1998 the National Marine Fisheries Service (NMFS) first listed the Central Valley steelhead as threatened under the Endangered Species Act (ESA). After several court challenges, NMFS issued a new final rule relisting the Central Valley steelhead on January 5, 2006 (71 FR 834). In a separate process regarding terms of the 1996 FERC license amendments for the Project, NMFS staff provided input to a draft limiting factors analysis for Tuolumne River salmonids (Mesick et al 2007) and included recommendations for developing abundance estimates, habitat use surveys and anadromy determination of resident *O. mykiss*. These recommendations were conceptually used to develop the Districts FERC Study Plan (TID/MID 2007) which was the subject of an April 3, 2008 FERC Order. As part of the Order, the Districts are required to conduct population estimate surveys in summer (June/July) and winter (February/March), starting in summer 2008 to determine *O. mykiss* population abundance by habitat type.

The purpose of the proposed *O. mykiss* population surveys is to provide population size estimates over several sampling seasons of differing environmental conditions to determine habitat use and needs within the lower Tuolumne River. The surveys will be used to examine the following hypotheses:

Hypothesis 1: Summertime distribution of suitable habitat by observed life stages of *O. mykiss* is related to ambient river water temperature.

Hypothesis 2: Habitat use by *O. mykiss* juveniles and adults observed in the Tuolumne River occurs at the same density in both restored and nearby reference sites.

As recommended by Stillwater Sciences (Stillwater), the surveys will employ a two-phase sampling approach of potential *O. mykiss* habitat using snorkel surveys for the development of a “bounded count” population estimate (Hankin and Mohr 2001). Although the methodology presented below discusses both repeated dive counts and calibration by depletion electrofishing, current ESA permit restrictions for both NMFS Section 10(a)(1)(A) permit No’s 1280 (TID) and 1282 (Stillwater) do not allow sufficient incidental take to conduct the second phase surveys at this time using electrofishing. Discussions with NMFS permitting staff and Stillwater have occurred since submittal of the 2007 FERC Study Plan, resulting in a pending formal request to NMFS by Stillwater for modification of Permit 1282 (see Section 6 below). The Section 10 Permit 1280 issued to TID in 2005 authorized only up to 5 juvenile *O. mykiss* annually by electrofishing that was further restricted to River Mile 25–30 during September to November. Thus that permit is not applicable or adequate to the season, location, and fish numbers needed to conduct the electrofishing for this population estimate study. Consequently, the July 2008 survey was conducted using snorkel surveys only as provided for in the 2007 study plan. It is not anticipated that the pending permit amendment request will be resolved prior to the winter 2009 survey, as such this will be conducted using snorkel surveys. If the pending amendment request is resolved prior to July 2008, then summer 2009 surveys will be conducted using the combined method presented below.

2 FIELD SAMPLING AND DATA COLLECTION

The two-phase stratified sampling design involves snorkeling pre-selected habitat units (e.g., riffle, run, pool, etc.) multiple times in order to quantify the variance associated with density and

subsequent population estimates. Habitat units are selected using stratified random sampling where the habitat types possess a pre-determined probability of occurrence within areas where *O. mykiss* have been frequently observed during the summer in the lower Tuolumne River, extending from approximately river mile (RM) 52–40 during summers and potentially extending to near the city of Waterford (RM 30) during colder winter conditions.

In a typical Phase 1 sampling approach, primary snorkel surveys (Edmundson et al. 1968, Hankin and Reeves 1998, McCain 1992, Dolloff et al. 1996) will be conducted across a subset of all habitat units. In Phase 2, approximately 20–70% of each habitat type sampled will be randomly selected for replicated surveys by either repeated dive counts or depletion electrofishing (Reynolds 1996). Although the bounded counts methodology was developed for use in smaller stream systems (Hankin and Mohr 2001), applying the methodology to a larger system such as the Tuolumne River is feasible provided key assumptions are satisfied. A critical assumption of the bounded counts approach is that all individuals have a chance of being observed. This may not be practically attainable due to the depths of some of the in-channel mining pits and also potentially due to low visibility conditions occurring at downstream locations or due to winter-time sediment inputs during rain events. Hankin and Mohr (2001) found that their survey designs were suitable for coho salmon (*O. kisutch*), but they were less confident about applying the methodology to *O. mykiss* juveniles because the fish’s furtive nature may violate the assumption that all fish have an observation probability >0. Sampling sites and methods may be modified following initial surveys because local conditions cannot be anticipated and may dictate the use of other schedules, locations, or techniques. Stillwater Sciences will notify TID, FERC, and permitting authorities if substantive changes in the study design, methods or schedule are anticipated.

2.1 Habitat Typing

On-the-ground mapping of potential habitat for *O. mykiss* will be delineated on digital ortho-rectified aerial photographs and information from previous habitat mapping efforts. Appendices A and B shows preliminary habitat units from RM 52–30 based upon habitat mapping conducted by Stillwater Sciences (2008) between La Grange Dam (RM 52) and Roberts Ferry Bridge (RM 40) (Appendix A) as well as preliminary habitat units from RM 40 to Waterford (RM 30) based upon mapping conducted by McBain & Trush (2004) and EA Engineering (1997) shown in Appendix B. The Appendix B habitat maps will be updated for flow and morphological characteristics in the field in late February and late June in each year. The final habitat maps will delineate all potential *O. mykiss* habitats according to the major types listed in Table 1, as well as transitional habitats that may be preferentially used by various size classes (i.e., pool heads, pool bodies, pool tails, run heads, run bodies, run tails, and riffles).

Table 1. Coarse scale habitat types to be used during snorkel surveys

Habitat Type	Description ^a	Approximate Depth
Riffle	Shallow with swift flowing, turbulent water. Partially exposed substrate dominated by cobble or boulder. Gradient moderate (less than 4%).	0–4 ft
Run	Fairly smooth water surface, low gradient, and few flow obstructions. Mean column velocity generally greater than one foot per second (fts ⁻¹).	4–10 ft
Pool	Slow flowing, tranquil water with mean column water velocity less than 1 fts ⁻¹ .	>10 ft

^aMajor habitat types determined based upon observed hydraulic conditions (McCain 1992, Thomas and Bovee 1993, Cannon and Kennedy 2003)

A Geographic Information System (GIS) will be used to update and refine habitat maps prior to thorough field verification of flow, depth, and habitat conditions in the river. Within each reach, individual habitat units will be digitized as two-dimensional features of varying shapes, or polygons, where each unit is a discrete functional habitat, as defined above. This approach is consistent with the general techniques of McCain (1992), Thomas and Bovee (1993), and Cannon and Kennedy (2003) and allows a flexible approach to evaluating habitat and habitat use patterns at a scale that can be easily delineated given available data, readily depicted, and is ecologically meaningful for aquatic species.

Habitat units will be assigned a natural sequence order (NSO), starting at one which is the first unit at the upstream end of the site, and a habitat type unit number (1...N pools, runs and riffles). The maximum depth, length and width (usually at 1/3 and 2/3 of the units length) will be recorded and flagging tied at both upstream and downstream ends of units to be surveyed. Pertinent information such as date, unit number, and type is included on the flag. Lastly, the upper and lower end of each unit will be located by GPS and mapping from previous efforts will be verified or updated.

2.2 Sample Site Selection

After all potential habitat units are typed and all pertinent information recorded, a subset of each habitat unit type will be selected for single-pass snorkel surveys. Although additional units may be selected at gravel augmentation and other in-channel restoration sites (See Hypothesis 2), selection for sampling proceeds by random selection of the starting sampling unit in the upper survey section, followed by a systematic uniform sampling of the remaining units in the survey reach. For example, every 3rd, 4th or larger selection interval will be used to distribute the selected units uniformly across the survey reach.

Because the total length of river sampled affects the confidence bounds of the resulting *O. mykiss* population estimates, at least 10% of the total length of a given habitat type and a minimum of 5 units of each type will be sampled. Based upon preliminary habitat mapping and median unit lengths of various habitat types, Table 2 shows that 63 sampling units for the winter surveys will be selected from representative locations between RM 52–30 to meet the minimums above. This estimate further assumes that, since detailed habitat type mapping has not been conducted from RM 40–30, habitat type distribution and median length from RM 40–30 are similar to RM 52–40, as determined by summer 2008 habitat type mapping (Stillwater Sciences 2008). The exact number sampled will be determined after random selection of the habitat units prior to study implementation.

During summer, an estimated 35 units will be selected for single-pass snorkel survey from representative locations between RM 52–40 (Table 2). For both winter and summer surveys, the number and location of habitat units may be adjusted if initial systematic sampling does not allow the study to adequately to test Hypothesis 2.

Table 2. Estimated number of sampling units that will meet study design assumption of sampling at least 10% of the total length of a given habitat type.

Habitat Type	Total length (ft) RM 52-40 ^a	Estimated total length (ft) RM 40-30 ^b	Estimated total length (ft) RM 52-30	Median length (ft) ^c	# of units to be sampled Winter 2009 RM 52-30 ^d	Estimated sampled Length Winter 2009	# of units to be sampled Summer 2009 RM 52-40 ^d	Estimated sampled Length Summer 2009
Riffle	14,320	13,590	27,910	322	9	10%	5	11%
Pool head	619	618	1,237	106	9	77%	5	86%
Pool body	6,741	6,795	13,536	393	9	26%	5	29%
Pool tail	781	618	1,399	124	9	80%	5	79%
Run head	2,067	1,853	3,920	51	9	12%	5	12%
Run body	37,350	35,829	73,179	843	9	10%	5	11%
Run tail	2,393	2,471	4,864	54	9	10%	5	11%
Total	64,271	61,775^e	126,046		63		35	

^aFrom Stillwater Sciences (2008)

^bAssumes same proportion of habitat types as from RM 52-40

^cAssumes median habitat unit lengths from RM52-40 are proportional to median lengths along RM 40-30.

^dAssumes at least 10% of the total length of each habitat type will be sampled; Estimates based upon 10% of the total length of a habitat type by median habitat unit length to determine a minimum number of units

^eActual river length from RM 40-30

2.3 Sampling Period

Winter sampling will begin in late February with systematic random selection of habitat units from RM 52-30, based upon summer 2008 maps (Appendix A) and previous habitat typing between RM 40-30 (Appendix B). Following habitat selection, Stillwater will use single-pass snorkel surveys and second phase calibration surveys within units of each type to develop uncertainty and bias estimates. Second phase sampling will be conducted using multi-pass snorkel surveys and/or depletion electrofishing methods as allowed under applicable permits (See Section 6).

Summer sampling will use habitat maps from RM 52-40 developed in summer 2008 (Appendix A). Although no additional habitat mapping is anticipated following winter 2009 surveys, habitat unit flagging will be established in advance of each snorkel survey effort and seasonal changes in habitat distribution may force revision of habitat type maps, specifically the upper and lower boundaries of habitat units and/or channel margins, prior to summer 2009 surveys.

2.4 Measurement Parameters and Sampling Methods

Multiple parameters will be measured in order to meet the objectives for this study (Table 3). Photos and GPS locations will be taken at each site, and site locations identified on GIS maps corresponding to mapped aquatic habitat units. General site information recorded at fish sampling locations will include site name, GPS coordinates, time, date, and crew member names. *In situ* water quality parameters (Temperature, dissolved oxygen, and conductivity) will be collected using a pre-calibrated multi-probe (YSI 85, Yellow Springs Instruments, Yellow Springs, OH). Underwater visibility will also be estimated into the sun and away from the sun using a Secchi disk to monitor any changes in visibility. Dissolved oxygen probes will be recalibrated at each site and checked for accuracy against concentrations measured in Winkler titrations (Grasshoff et al 1983) at the beginning and end of the sampling effort using a dissolved oxygen test kit.

Table 3. Measurement parameters and methods for snorkel surveys

Parameter	Method	Metric/Descriptor	Method Reporting Limit
<i>Habitat Typing Attributes</i>			
Natural sequence order (Reach ID – Habitat unit #)	N/A	A-1, A-2, A-3, ...	N/A
Latitude/Longitude	Handheld GPS receiver	UTM	N/A
Habitat type	Visual estimation	See Table 1	N/A
Average unit width	Horizontal distance	meters (feet) (measured at multiple transects)	3 ft (1 m)
Average unit length	Horizontal distance	meters (feet)	3 ft (1 m)
Maximum/minimum depth	Vertical distance	meters (feet)	1 ft (0.3 m)
Bed substrate composition	Visual estimation	bedrock, boulder, cobble, gravel, organic, sand, silt	10%
Cover type	Visual estimation	none, boulder, cobble, IWM, bedrock ledges, overhead vegetation, aquatic vegetation	10%
<i>Field Data During Snorkel Surveys</i>			
Temperature	EPA 170.1	°C	0.1 °C
Dissolved Oxygen	SM 4500-O	mg/L	0.0 mg/L
Conductivity	SM 2510A	umhos/cm	1.0 umhos/cm
Visibility	Secchi depth	meters (feet)	0.01 m (0.1 ft)
Date/Start time/End time	N/A	Day/month/year	N/A
Number of Individuals	Visual estimation	Number	1
Fish length – snorkeling	Visual estimation	millimeter	50 mm
Fish length – electrofishing	Fork length	millimeter	1 mm
Weight - electrofishing	Electronic balance	gram	0.1 g

2.4.1 Snorkel Surveys

Snorkel surveys will be conducted during daylight hours (7:00am–5:00pm winter; 6:00am–8:00pm summer). A two phase survey design will be used to survey the seven different strata (Table 4). At the first phase, single-pass dive surveys will be conducted by a four to five person crew depending upon river flows and underwater visibility. Sampling units will generally be sampled from downstream to upstream in dive lanes using a zigzag pattern, passing fish and allowing them to escape downstream of the diver. If fish are observed to escape upstream, the diver will take care to avoid counting these fish twice. Divers will record their observations of pertinent attributes (Table 3) and numbers of *O. mykiss* and Chinook salmon (*O. tshawtscha*) observed; with fish lengths to be estimated in 50 mm size ranges using a scale model or markings on the slates to correct for underwater size distortion. After the first dive pass is completed a tab is then pulled to determine if the unit is included in the second phase of sampling.

Table 4. Preliminary sample unit selection and survey count.

Habitat	Winter 2009				Summer 2009			
	Phase I Dives		Phase II Survey		Phase I Dives		Phase II Survey	
	Initial Units	Passes	Repeat Units	Passes	Initial Units	Passes	Repeat Units	Passes
Riffle	9	1	2	2	5	1	2	2
Pool head	9	1	2	2	5	1	2	2
Pool body	9	1	2	2	5	1	2	2
Pool tail	9	1	2	2	5	1	2	2
Run head	9	1	2	2	5	1	2	2
Run body	9	1	2	2	5	1	2	2
Run tail	9	1	2	2	5	1	2	2
	Total	63	Total	28	Total	35	Total	28

The second phase of sampling collects data that will later be used to extrapolate dive counts to total population estimates by three passes of either repeated dive counts or depletion electrofishing. Ideally, if the count of *O. mykiss* from the Phase 1 snorkel survey is less than or equal to 20 individuals then three additional dive passes are made. If electrofishing is permitted, all units with a count of juvenile *O. mykiss* counts greater than 20 individuals will be surveyed by electrofishing. Lastly, occurrence of other native and non-native fish species will be recorded as presence/absence.

2.4.2 Electrofishing at Riverine Sites

If employed during the summer 2009 survey, electrofishing will be conducted by a 4 person crew during the daylight hours (6:00am-8pm) following the dive surveys. Ideally, 3-pass electrofishing will be used on all second phase dive units where the first dive pass exceeded 20 *O. mykiss*. Dive units that require electrofishing for dive calibration will be completed as soon as possible after the dive survey.

Shallow water habitat may be sampled using back pack electrofishing units while deep water habitat may be sampled using a boat electrofishing unit. Back pack electrofishing in shallow waters less than 3–4 ft depth will be conducted using two or more Smith-Root back pack electrofishers (Model LR-24 or Model 12 with 11-inch anode rings and standard “rat-tail” cathodes). Boat electrofishing may be used in deeper riverine habitats using a boat mounted Smith Root 1.5 KVA electrofishing unit. To ensure the health of all fish captured during electrofishing, all electrofishing will be conducted in accordance with NMFS (2000) electrofishing guidelines and an electrofishing logbook will be maintained and updated at each sampling site.

Depending upon river flows and depth, electrofishing will use block nets placed at the upstream and downstream ends of the unit to be fished, taking care to avoid disturbance of the unit during net set-up. Block nets will be set up where possible to prevent fish from moving out of the unit. If block nets are not feasible, then a snorkeler may be stationed at the upstream end of a unit to observe any fish moving out of the unit.

First pass electrofishing will proceed slowly and deliberately upstream from the downstream end of the unit; members of an electrofishing crew will move to the top and back down to the bottom working closely together. To maintain equal effort on subsequent passes, electrofishing time (seconds) will be recorded to allow for any adjustments in sampling effort. A fourth pass will be conducted if one of the following applies:

1. The number of *O. mykiss* caught on the 2nd pass exceeds the number of *O. mykiss* caught on the 1st pass.
2. The number of *O. mykiss* caught on the 3rd pass is greater than or equal to 25 percent of number caught on the 2nd pass.

The procedure may be modified in riffle habitats to facilitate capture of shocked fish in fast water. In the riffle strata, a pass consists of a sweep from the top to the bottom of the unit. Depending on the water velocity, block nets may or may not be set at the upstream end of riffle units.

2.4.3 Fish Handling Protocols

Any fish captured during electrofishing surveys will be processed, and information collected regarding species identification, fork length (FL, mm), weight (g), and, if applicable, notes on general condition. All fish will be rapidly retrieved using dip nets and placed immediately into aerated live wells or buckets with water. Large fish will be kept separate from juvenile fish to avoid confinement predation. Fish will be identified to species and origin (hatchery or wild stock) where possible. Fish that are weighed and measured will be anesthetized using clove oil to minimize handling stress. After all fish are identified, counted, and measured, fish will be held for approximately 10 minutes, until they show signs of “normal” swimming patterns and behavior.

2.5 Hypothesis Testing

The purpose of the proposed *O. mykiss* population surveys is to provide population size estimates over several sampling seasons of differing environmental conditions to determine habitat use and needs within the lower Tuolumne River. The surveys will be used to examine the following hypotheses:

Hypothesis 1: Summertime distribution of suitable habitat by observed life stages of *O. mykiss* is related to ambient river water temperature.

Hypothesis 2: Habitat use by *O. mykiss* juveniles and adults observed in the Tuolumne River occurs at the same density in both restored and nearby reference sites.

While the selection for sampling proceeds by random selection of the starting sampling unit in the upper survey section, followed by a systematic uniform sampling of the remaining units in the survey reach, additional units adjacent to or near restoration sites may be non-randomly selected to provide treatment and control locations to test Hypothesis 2, especially during winter 2009 surveys when low ambient river water temperatures obviate the need to test Hypothesis 1.

2.6 Field Work Notification

To ensure field staff safety and to satisfy scientific collecting permit requirements, the parties listed in Table 5 will be notified in advance of the proposed sampling in as required to confirm sampling dates.

Table 5. Field Work Notification

Contact	Affiliation	Address	Phone and Email
Tim Ford	TID	333 East Canal Dr. Turlock, CA 95380	209.883.8275 tjford@tid.org
Tim Heyne	CDFG	P.O. Box 10 La Grange, CA 95329	209.853.2533 x1# theyne@dfg.ca.gov
Jeffery Jahn	NMFS	777 Sonoma Ave. Rm 325 Santa Rosa, CA 95404	707.575.6097 Jeffrey.Jahn@noaa.gov

Prior to mobilization, planned river operations by the Districts will be checked to determine if fish sampling would be safe under the anticipated flow and all parties will be notified of any delay or modification to the sampling schedule.

3 QUALITY ASSURANCE

The objective of data collection for this Project is to produce data that represent as closely as possible, *in situ* conditions of the Tuolumne River with respect to river flow conditions, water quality, abundance and habitat use by *O. mykiss*. To meet this objective, field sampling, sample preparation, and analysis will follow general guidelines outlined in USEPA (2002) by ensuring that:

- the project's objectives, hypotheses and data quality objectives are identified and agreed upon,
- the intended measurements and methods are consistent with project objectives,
- the assessment procedures are sufficient for determining if data of the type and quality needed and expected are obtained, and
- any potential limitations on the use of the data can be identified and documented.

Aquatic environments are inherently variable, but management decisions must be based on a data from a limited number of locations and often collected in short time periods. How well the information collected represent the reach or river-wide fish population depends upon a systematic approach to quality assurance.

3.1 Data Quality Objectives for Measurement Data

The data quality parameters used to assess the acceptability of the data are precision, accuracy, representativeness, comparability, and completeness. Precision measures the reproducibility of measurements under a given set of conditions. Analytical precision is limited to water quality and physical habitat characteristics (Table 6). Accuracy is an expression of the degree to which a measured or computed value represents the true value. Field accuracy is controlled by adherence to sample collection procedures.

Table 6. Data quality objectives for field parameters

Parameter	Units	Accuracy	Precision	Completeness
Dissolved Oxygen	mg/L	± 0.5	10%	90%
Temperature	°C	± 0.5	5%	90%
Conductivity	umhos/cm	± 5%	± 5%	90%
Depth	meters	± 0.2	N/A	N/A
Visibility (Secchi)	meters	± 0.05	N/A	N/A

- Representativeness expresses the degree to which data accurately and precisely represent an environmental condition. For this study, monitoring site selection will be conducted based on physical habitat attributes. Additionally, specific measurement parameters have been identified as relevant based on numerous studies indicating factors associated with species distribution.
- Comparability expresses the confidence with which one data set can be evaluated in relation to another data set. For this biological assessment, comparability of data will be established through the use of standard analytical methodologies and reporting formats.
- The project goal for completeness, a measure of the amount of data that is determined to be valid in proportion to the amount of data collected, will be 90% for analytical water quality parameters. The data quality objective for completeness for all components of this study is 90%.

3.2 Training Requirements/Certification

Specialized training is required for the proposed sampling activities, however none of the sampling activities require outside certification from an agency or another entity. Required permits for biological sampling are discussed in Section 5. Field crews will be staffed by a variety of qualified personnel, which due to the nature of extended field activities, will necessarily be rotated in and out of the field.

3.3 Instrument/Equipment Testing, Inspection, and Maintenance Requirements

To ensure proper equipment performance in the field, maintenance and operational procedures, including preventative maintenance, will be performed on all YSI multiprobes (temperature, dissolved oxygen, and conductivity). YSI maintenance will be recorded in a logbook with the date the maintenance was performed and the initials of the technician. When the instruments are not deployed, the calibration or storage cup will be used to protect sensors from damage and desiccation.

3.4 Instrument Calibration and Frequency

Field probes used for field sampling will be calibrated prior to use, midway through each sampling event, and at the end of each sampling event. Measurement devices for conductivity will be checked against a standard whose source is different than that selected for calibration. Dissolved oxygen will be checked against aerated water whose oxygen content is established by the Winkler method (Grashoff et al 1983). Temperature does not require calibration because of the unvarying nature of the temperature sensor and its conditioning circuitry.

3.5 Reconciliation with Data Quality Objectives

If data do not meet the project's specifications, the following actions will be taken. First, the task leaders working with the field crew leaders (in some cases they will be the same person) will review the errors and determine if the problem is equipment failure, calibration/maintenance techniques, or monitoring/sampling techniques. They will suggest corrective action. If the problem cannot be corrected by training, revision of techniques, or replacement of supplies/equipment, then the task leaders will review the data quality objectives (DQOs) and determine if the DQOs are feasible. If the

specific DQOs are not achievable, they will determine whether the specific DQO can be relaxed, or if the parameter should be eliminated from the monitoring program.

3.6 Data Management

All field data will be amassed in a quality-checked database and summarized. QA checks will be applied to all data before data entry and data will be stored on Stillwater Sciences servers. Full backup of data from all offices is done on a weekly basis, while differential backup (files that have changed since the last full backup) is done on a nightly basis. The backup process is accomplished with a Fast Tape Library and backup processes are completed during off-peak hours. Two sets of tapes are taken offsite by two Information Technology (IT) staff members on a weekly basis to ensure recovery in case of failure or catastrophe.

4 DATA ANALYSIS

Data analysis will be conducted to summarize *in situ* water quality and fish counts in each sampling strata. Bounded counts or depletion estimators will be used to determine populations and linear density for each sampled unit, together with estimates of uncertainty. In addition to comparisons of fish density between sampling strata, the density estimates and uncertainties will be propagated across the unsampled areas for an overall population estimate. Exploratory multiple regression analysis will also be used to determine relationships between fish density and recorded habitat variables.

5 REPORTING

A data report will be prepared for use with permitting authorities that includes: date, time, and location of sampling activities; species and number of species collected; and a copy of field data sheets. Results of the winter 2009 surveys will be transmitted to TID electronically within three weeks of the survey completion (April/May 2009). A client review draft of the technical report covering the results of both winter and summer 2009 surveys will be submitted to TID by August 24, 2009. Assuming an internal and Agency review comments are received within one and three weeks of issuance of the client review and Agency review drafts, respectively, the Agency review draft will be available by September 8, 2009 and final report will be complete by October 16, 2009.

6 PERMITTING REQUIREMENTS

Stillwater Sciences will maintain the following permits to sample fish populations that may be present:

- NMFS Section 10(a)(1)(A) permit 1282
- California Department of Fish and Game individual Scientific Collection Permits.

A NMFS Section 10(a)(1)(A) permit 1282 has been obtained and all NMFS guidelines (e.g., notification, data gathering, preservation) will be followed if any Central Valley steelhead are captured. Under that existing NMFS permit, electrofishing is limited to an authorized incidental take of 40 juvenile *O. mykiss* and the <5% unintentional mortality limit, and no adults. An amendment to the sampling description was submitted to NMFS on June 2, 2008 with increased take limits for handling electrofishing of 100 adults and 200 juveniles at an unintentional mortality rate of <10%. Mr. Jeffrey Jahn of NMFS will be notified at least two weeks prior to applicable sampling to confirm

sampling dates and locations. Electrofishing under an amended permit will be suspended in the event that the authorized incidental take limits were exceeded and all subsequent calibration surveys would be made by repeat dive surveys. Annual reporting will be provided to Mr. Jeffrey Jahn of NMFS by March 1, of each year.

CDFG Scientific Collecting Permits (SCPs) will be maintained for species potentially present in the project area. CDFG guidelines (e.g., notification, data gathering, and preservation) will be followed if special-status species are captured and the CDFG 24-hr dispatch (916.446.0045) will be notified should unrelated events result in fish kills.

No intentional mortality or removal of special-status species from the wild is included in this study plan. In the event unintentional mortality occurs beyond the take permit limits, NMFS staff will be contacted within 24 hrs and a fin-clip will be provided to the Salmonid Genetic Repository. CDFG will also be contacted to determine the disposition of the individual specimen and whether the individual may be retained for otolith analysis.

7 REFERENCES

- Cannon, T.C., and T. Kennedy. 2003. Snorkel survey of the lower American River 2003. Prepared by Fishery Foundation of California, San Francisco for U. S. Fish and Wildlife Service, Central Valley Project Improvement Program, Sacramento, California.
- Dolloff, C. A., J. Kershner, and R. Thurow. 1996. Underwater observation. Pages 533–554 in B. R. Murphy and D. W. Willis, editors. Fisheries techniques, 2nd edition. American Fisheries Society, Bethesda, Maryland.
- EA Engineering. 1997. Tuolumne River GIS Database Report and Map. Prepared for the Turlock Irrigation District and Modesto Irrigation District by EA Engineering. FERC Project No. 2299, 1996 FERC Report, Volume VII, Attachment 96-14.
- Edmundson, E.F., E. Everest, and DW. Chapman. 1968. Permanence of station in juvenile chinook salmon and steelhead trout. *Journal of the Fisheries Research Board of Canada* 25(7): 1453–1464.
- Grashoff, K., M. Erhardt, and K. Kremling. 1983. *Methods in Seawater Analysis*. 2nd ed. Verlag Chemie, Weinheim.
- Hankin, D.G. and M. Mohr. 2001. Improved Two-Phase Survey Designs for Estimation of Fish Abundance in Small Streams. Preprint from David G. Hankin, Department of Fisheries Biology, Humboldt State University, Arcata, CA
- Hankin, D.G., and G.H. Reeves. 1988. Estimating total fish abundance and habitat area in small streams based on visual estimation methods. *Can. J. Fish. and Aqu. Sci.* 45:834-844.
- McCain, M.E. 1992. Comparison of habitat use and availability for juvenile fall chinook salmon in a tributary of the Smith River, California. *FHR Currents*. No. 7. USDA Forest Service, Region 5.
- McBain & Trush. 2004. Habitat Maps for the upper 15.8 miles of the Gravel Bedded Reach Appendix D *In* Coarse Sediment Management Plan for the Lower Tuolumne River. Revised Final. Prepared for Tuolumne River Technical Advisory Committee, Turlock and Modesto Irrigation Districts, USFWS Anadromous Fish Restoration Program and the CALFED Bay Delta Authority. Prepared by McBain & Trush, Arcata, CA July 20.
- Mesick, C., J. McClain, D. Marston, and T. Heyne. 2007. Draft Limiting Factor Analyses & Recommended Studies for Fall-run Chinook salmon and Rainbow Trout in the Tuolumne River, Attachment 2 to USFWS comments on FERC study plan. Available at the FERC website: http://elibrary.FERC.gov/idmws/file_list.asp?accession_num=20070314-0089
- Stillwater Sciences. 2008. July 2008 Population Size Estimate of *Oncorhynchus mykiss* in the Lower Tuolumne River. Prepared for the Turlock Irrigation District and Modesto Irrigation District by Stillwater Sciences, Berkeley, CA. October 2008.
- TID/MID (Turlock and Modesto Irrigation Districts). 2005. Ten Year Summary Report of Turlock Irrigation District and Modesto Irrigation District Pursuant to Article 58 of the License for the Don Pedro Project, No. 2299. 1 Volume. March.

TID/MID. 2007. Tuolumne River Fisheries Study Plan - Don Pedro Hydroelectric Project (FERC NO. 2299). Prepared by T. Ford, N. Hume, S. Wilcox, and R. Yoshiyama for Turlock Irrigation District and Modesto Irrigation District. July 13. Available at the FERC website:
http://elibrary.FERC.gov/idmws/file_list.asp?accession_num=20070718-0082

Thomas, J.A., and K.D. Bovee. 1993. Application and testing of a procedure to evaluate transferability of habitat suitability criteria. *Regulated Rivers: Research and Management* 8: 285 - 294.

USEPA (U.S. Environmental Protection Agency). 2002. Guidance for Quality Assurance Project Plans EPA QA/G-5. EPA/240/R-02/009. U.S. Environmental Protection Agency, Office of Environmental Information, Washington, DC. December.

Appendix B: 2008 Habitat Maps




Tuolumne River - O. mykiss BCE Surveys, 2008-2011





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 02





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 03





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 04





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 05





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 06





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 07





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □

- Tile Boundary (shown white on the map)
- River Miles

METADATA

Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 08





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 09





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles


METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles ●


METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs █
- Riffles
- Pools
- Tile Boundary (shown white on the map)
- River Miles


METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA


Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA

Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs █
- Riffles █
- Pools █
- Tile Boundary (shown white on the map)
- River Miles


METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.


Tile 18



FOT BOBCAT FLAT RESTORATION SITE



Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles ●

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)


Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 19





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles ●

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 20





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 21





Tuolumne River - O. mykiss BCE Surveys, 2008-2011




- Runs █
- Riffles █
- Pools █
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)


Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 22





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA

Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs █
- Riffles █
- Pools █
- Tile Boundary (shown white on the map)
- River Miles

METADATA

Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs █
- Riffles █
- Pools █
- Tile Boundary (shown white on the map)
- River Miles

METADATA
 Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)


Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.

Tile 25





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA


Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □
- Tile Boundary (shown white on the map)
- River Miles

METADATA

Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.





Tuolumne River - O. mykiss BCE Surveys, 2008-2011



- Runs □
- Riffles □
- Pools □

- Tile Boundary (shown white on the map)
- River Miles

METADATA



Tiles 29 to 47: NAIP, 6/29/2009 (130 cfs)
 Tiles 1 to 29: Sanborn imagery, 09/25/2005 (335 cfs)

Wetted perimeter were first based on EA_mapping data (90's) at 230 cfs, and later refined using 2005 & 2009 NAIP and field measurements from 2008 and 2009 surveys to adjust for channel migration.


Tile 28





Appendix C: 2004 Habitat Maps




TUOLUMNE RIVER HABITAT MAP SHEET INDEX












- Legend
- Proposed coarse sediment introduction site
 - Area with sensitive *O. mykiss* habitat
 - Spawning area (@ 300 cfs)
 - Riffle
 - Run
 - Special run pool
 - Pool
 - Glide
 - Medial or lateral bar
 - Submerged lateral bar
 - Deep backwater
 - Shallow backwater
 - Island complex
 - Off-channel pond
 - Bedrock
 - Access road
 - Cross section
 - Station alignment









Legend


- Proposed coarse sediment introduction site
- Area with sensitive *O. mykiss* habitat
- Spawning area (@ 300 cfs)
- Riffle
- Run
- Special run pool
- Pool
- Glide
- Medial or lateral bar
- Submerged lateral bar
- Deep backwater
- Shallow backwater
- Island complex
- Off-channel pond
- Bedrock
- Access road
- Cross section
- Station alignment
















Appendix D: Habitat Data

Table D-1. Physical habitat types and dimensions of surveyed areas in the lower Tuolumne River (RM 52-40).

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
1	51.8		Yes	140	75	10,537	5.0	8.0	Pool head
2	51.7			450	143	64,161	18.0	28.0	Pool body
3	51.7			157	61	9,600	1.5	3.0	Pool tail
4	51.6	Yes	Yes	85	124	10,506	3.0	5.0	Pool head
5	51.6	Yes	Yes	393	129	50,702	18.0	25.0	Pool body
6	51.5			250	89	22,309	4.0	6.0	Pool tail
7	51.5			292	68	19,851	3.0	6.0	Riffle
8	51.4			117	82	9,562	5.0	6.0	Run head
9	51.1			2047	97	199,103	6.0	8.0	Run body
10	51.0			182	86	15,733	3.5	4.5	Run tail
11	50.9	Yes		457	99	45,397	10.0	16.0	Pool body
12	50.8	Yes	Yes	843	128	107,699	4.0	7.0	Run body
13	50.8			93	86	7,988	1.5	3.0	Run tail
14	50.6		Yes	708	65	45,670	1.5		Riffle
15	50.6	Yes		161	85	13,760	6.0	7.0	Run head
16	50.5	Yes		704	132	92,609	5.0	8.0	Run body
17	50.4			59	146	8,600	2.5	3.0	Run tail
18	50.3	Yes		941	130	121,948	1.5	2.0	Riffle
19	50.3	Yes	Yes	59	109	7,193	4.0	8.0	Run head
20	50.1	Yes		848	151	107,630	3.0	4.0	Run body
21	50.1			70	119	8,333	1.5	2.0	Run tail
22	50.1	Yes		132	127	16,750	1.0	1.5	Riffle
23	50.0			93	133	12,379	4.0	6.0	Run head
24	49.9		Yes	1007	199	200,462	4.0	8.0	Run body
25	49.8			274	154	42,115	2.0	4.0	Run tail
26	49.7	Yes		527	139	72,991	1.5	2.0	Riffle
27	49.7	Yes	Yes	127	86	10,955	4.0	6.0	Pool head
28	49.6	Yes	Yes	161	89	14,345	6.0	9.0	Pool body
29	49.6			112	85	9,490	1.5	2.5	Pool tail
30	49.6			50	110	5,520	3.0	5.0	Run head
31	49.3			1440	115	166,115	2.5	3.5	Run body
32	49.3			132	137	18,071	2.0	2.5	Run tail
33	49.2			552	126	69,509	1.5	2.5	Riffle
34	49.2			112	65	7,283	2.0	3.0	Run head
35	49.1			321	82	26,475	3.0	5.0	Run body
36	49.1			44	103	4,532	1.5	2.0	Run tail
37	49.1			78	97	7,594	1.5	2.0	Riffle
38	49.1		Yes	43	83	3,559	2.0	3.5	Run head
39	49.1			240	81	19,424	2.5	4.0	Run body
40	49.0			23	95	2,180	2.5	3.0	Run tail
41	48.8			1080	114	122,953	1.5	3.0	Riffle

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
42	48.8	Yes		36	97	3,505	1.5	2.0	Run head
43	48.7	Yes		749	93	69,528	2.5	4.0	Run body
44	48.7			39	110	4,304	2.0	3.0	Run tail
45	48.4		Yes	1275	117	149,495	1.5	2.0	Riffle
46	48.4			92	102	9,378	1.5	2.0	Run head
47	48.3			915	111	101,397	3.5	5.0	Run body
48	48.2			153	127	19,368	1.5	2.0	Run tail
49	48.2			346	75	25,887	1.5	2.0	Riffle
50	48.2			40	60	2,392	2.0	2.0	Run head
51	48.1		Yes	380	53	20,027	5.0	8.0	Run body
52	48.1			114	56	6,430	3.0	3.5	Run tail
53	48.0		Yes	234	54	12,554	1.5	2.0	Riffle
54	48.0	Yes	Yes	164	89	14,569	5.0	7.0	Pool head
55	47.2			4036	143	579,150	7.0	15.0	Pool body
56	47.2			136	115	15,575	1.5	2.5	Pool tail
57	47.1			740	80	58,852	1.5	2.0	Riffle
58	47.0			136	85	11,535	2.0	3.0	Run head
59	46.9			472	76	36,067	4.0	6.0	Run body
60	46.9			137	86	11,760	1.5	2.5	Run tail
61	46.9			318	81	25,666	1.0	2.0	Riffle
62	46.9		Yes	64	85	5,428	1.5	2.0	Run head
63	46.8			188	90	16,848	2.0	3.0	Run body
64	46.8			126	131	16,480	1.0	2.5	Run tail
65	46.8			100	123	12,268	0.8	1.5	Riffle
66	46.8			153	96	14,675	1.5	2.0	Run head
67	46.0			3829	97	370,148	4.0	6.0	Run body
68	46.0			89	133	11,835	1.5	2.0	Run tail
69	45.9			234	95	22,286	4.0	7.0	Run body
70	45.9	Yes		277	76	21,181	1.5	2.0	Riffle
71	45.9			61	93	5,701	2.0		Run head
72	45.8			243	94	22,751	2.5	3.5	Run body
73	45.8			125	64	7,976	1.5	2.0	Run tail
74	45.7			243	40	9,820	0.8	1.8	Riffle
75	45.7			90	35	3,141	1.5	2.0	Run head
76	45.7			88	50	4,433	1.5	4.0	Run body
77	45.7			32	99	3,153	1.5	2.0	Run tail
78	45.6			675	109	73,797	1.5	2.0	Riffle
79	45.6			85	178	15,127	1.5	2.0	Run head
80	45.4			1040	120	124,357	3.5	5.0	Run body
81	45.3		Yes	301	101	30,519	7.0	11.0	Pool body
82	45.3			126	220	27,658	2.0	3.0	Run head
83	45.1		Yes	1182	97	114,144	4.0	6.0	Run body
84	45.1			94	113	10,640	1.5	5.0	Run tail
85	45.0			394	52	20,673	1.5	2.0	Riffle

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
86	45.0	Yes	Yes	53	41	2,181	2.0	3.0	Pool head
87	44.9			101	71	7,213	5.0	8.0	Pool body
88	44.9			80	121	9,661	3.0	4.0	Pool tail
89	44.8			734	59	43,114	1.5	2.5	Riffle
90	44.8	Yes	Yes	22	107	2,350	0.8	1.5	Run head
91	44.8			318	62	19,745	1.5	2.5	Run body
92	44.8			15	25	368	1.0	1.5	Run tail
93	44.7	Yes		100	30	3,032	1.5	2.0	Riffle
94	44.7			47	26	1,217	1.0	1.5	Run head
95	44.7			248	67	16,708	4.0	8.0	Run body
96	44.7			34	87	2,950	1.5	2.0	Run tail
97	44.6			417	52	21,741	1.5	2.5	Riffle
98	44.6			20	49	984	2.0	2.5	Run head
99	44.6			203	53	10,740	3.0	4.0	Run body
100	44.5			20	59	1,182	1.0	1.5	Run tail
101	44.5	Yes	Yes	472	59	27,744	1.5	2.0	Riffle
102	44.5			10	68	681	2.0	2.5	Run head
103	43.9			3209	82	261,993	3.0	3.0	Run body
104	43.7	Yes	Yes	683	144	98,065	6.0	15.0	Pool body
105	43.3			2173	146	316,376	4.0	6.0	Run body
106	43.3			50	110	5,487	1.5	2.0	Run tail
107	43.2		Yes	326	81	26,534	1.5	2.0	Riffle
108	43.2			41	74	3,020	1.0	2.0	Run head
109	43.1			906	62	56,464	2.5	6.0	Run body
110	43.1			36	49	1,771	2.0	2.5	Run tail
111	43.0	Yes		238	42	10,077	0.8	1.2	Riffle
112	43.0	Yes		50	48	2,392	1.5	2.5	Pool head
113	43.0	Yes		159	166	26,397	5.0	7.0	Pool body
114	43.0			46	169	7,767	1.5	5.0	Pool tail
115	43.0			33	154	5,097	2.0	3.0	Run head
116	42.9	Yes		309	124	38,258	4.0	10.0	Run body
117	42.9			18	84	1,518	1.0	1.5	Run tail
118	42.9			77	57	4,403	1.0	2.0	Riffle
119	42.9	Yes		31	45	1,395	2.0	2.5	Run head
120	42.7			978	87	84,726	1.0	8.0	Run body
121	42.7			12	78	932	1.5	2.5	Run tail
122	42.7			89	48	4,288	1.0	3.0	Riffle
123	42.7		Yes	18	55	991	2.5	3.0	Run head
124	42.4		Yes	1571	77	120,609	2.0	5.0	Run body
125	42.4			69	96	6,600	1.5	2.0	Run body
126	42.3	Yes		227	55	12,478	1.0	3.0	Riffle
127	42.3			84	23	1,953	1.5	4.0	Run body
128	42.3			265	32	8,417	1.5	2.3	Riffle
129	42.2			25	28	699	1.5	3.0	Run head

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
130	42.1			1066	62	65,871	2.0	4.0	Run body
131	42.0			53	60	3,196	1.0	1.5	Run tail
132	41.9			521	64	33,202	1.0	1.5	Riffle
133	41.9	Yes		41	46	1,877	2.0	2.5	Run head
134	41.8	Yes		940	82	77,063	2.0	4.0	Run body
135	41.8			47	96	4,525	0.8	1.5	Run tail
136	41.7			300	90	27,080	0.8	1.5	Riffle
137	41.7			59	70	4,133	1.5	2.0	Run head
138	41.2			2512	123	308,848	3.0	6.0	Run body
139	41.2			125	151	18,858	1.0	1.3	Run tail
140	41.1			312	107	33,422	1.0	1.5	Riffle
141	41.1			102	163	16,604	1.5	2.0	Run head
142	41.0			666	185	122,933	2.0	4.5	Run body
143	41.0			83	182	15,121	0.8	1.3	Run tail
144	40.9			189	32	6,116	0.8	1.5	Riffle
145	40.9			62	39	2,425	1.5	2.0	Run head
146	40.5			2207	101	223,893	5.0	9.0	Run body
147	40.5			54	53	2,861	1.5	2.0	Run tail
148	40.4			638	53	33,978	1.5	2.5	Riffle
149	40.4			37	83	3,076	1.5	2.0	Run head
150	40.3		Yes	502	94	47,268	2.5	4.0	Run body
151	40.3			34	81	2,767	1.0	1.5	Run tail
152	40.2			503	53	26,860	0.8	1.5	Riffle
153	40.2			51	68	3,462	1.5	2.0	Run head
154	39.7			2569	123	317,216	3.0	7.0	Run body
155	39.7			26	142	3,699	1.5		Run tail
156	39.7		Yes	219	91	19,859	0.8	1.0	Riffle
157	39.6		Yes	86	62	5,294	3.0	4.0	Run head
158	39.5			857	97	82,763	6.0	6.6	Run body
159	39.5			98	81	7,993	2.5	3.0	Run tail
160	39.4			84	62	5,246	1.0	1.5	Riffle
161	39.4			123	41	5,102	3.5	4.5	Run head
162	39.3			713	50	35,662	5.0	7.5	Run body
163	39.3			151	80	12,041	3.5	5.0	Run tail
164	39.2			104	98	10,131	1.0	1.5	Riffle
165	39.2	Yes	Yes	93	117	10,818	3.5	4.0	Pool head
166	38.9	Yes	Yes	1496	90	134,259	6.5	9.9	Pool body
167	38.9			99	91	9,033	3.0	4.0	Pool tail
168	38.9	Yes	Yes	73	92	6,682	1.5	3.0	Riffle
169	38.9			76	108	8,227	4.0	5.0	Run head
170	38.8			498	77	38,331	5.5	7.2	Run body
171	38.8		Yes	121	83	10,096	7.0	10.5	Pool body
172	38.8	Yes		87	98	8,506	3.0	4.0	Run head
173	38.7	Yes		324	85	27,545	4.0	5.0	Run body

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
174	38.7			99	100	9,935	3.0	4.0	Run tail
175	38.7			61	118	7,163	1.5	2.3	Riffle
176	38.6			148	105	15,607	2.5	3.5	Run head
177	38.6			219	91	19,976	4.0	4.8	Run body
178	38.6			115	57	6,513	2.0	2.5	Run tail
179	38.5	Yes		412	55	22,840	1.2	2.0	Riffle
180	38.5			75	68	5,113	4.0	6.0	Run head
181	38.4			657	39	25,600	4.0	5.0	Run body
182	38.3			205	68	13,869	8.5	10.5	Pool body
183	38.3			183	66	12,189	4.5	10.5	Pool tail
184	38.3			129	102	13,154	2.5	6.0	Run head
185	38.2			137	139	18,966	2.0	2.5	Run body
186	38.2			134	149	19,976	2.0	2.0	Run tail
187	38.2			285	143	40,886	1.0	1.5	Riffle
188	38.1			86	93	7,964	2.5	4.0	Pool head
189	38.1			235	81	19,027	6.0	10.0	Pool body
190	38.1			55	145	7,947	2.5	4.0	Pool tail
191	38.1			89	115	10,283	1.0	2.0	Riffle
192	38.1			46	89	4,147	4.0	6.0	Pool head
193	38.0			378	83	31,490	8.0	13.0	Pool body
194	38.0			81	91	7,365	2.0	3.5	Pool tail
195	38.0			63	64	4,010	3.0	3.5	Run head
196	37.9			271	72	19,591	4.0	5.5	Run body
197	37.9			84	92	7,736	3.0	3.5	Run tail
198	37.8			227	75	17,099	2.0	2.5	Riffle
199	37.8			115	42	4,779	4.0	4.5	Pool head
200	37.7			926	78	72,513	4.0	6.6	Pool body
201	37.6			114	117	13,311	3.0	4.0	Pool tail
202	37.6			163	97	15,857	0.8	1.5	Riffle
203	37.6			130	88	11,423	2.0	3.0	Run head
204	37.5			618	91	55,953	2.5	3.5	Run body
205	37.4			102	77	7,851	2.0	3.0	Run tail
206	37.3			769	50	38,658	1.7	2.5	Riffle
207	37.3			99	58	5,710	2.5	4.0	Run head
208	37.1			916	57	51,803	3.5	4.5	Run body
209	37.1			58	52	3,054	2.0	3.0	Run tail
210	37.0			266	40	10,767	1.5	2.0	Riffle
211	37.0			127	36	4,530	5.0	7.0	Run head
212	36.9			370	80	29,741	5.5	7.6	Run body
213	36.9			85	98	8,321	2.0	3.0	Run tail
214	36.9			70	83	5,779	3.0	5.0	Pool head
215	36.9			126	58	7,330	7.0	10.5	Pool body
216	36.9			94	48	4,471	4.0	5.0	Pool tail
217	36.8			357	60	21,436	1.5	2.0	Riffle

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
218	36.8			157	75	11,815	3.0	4.0	Run head
219	36.6			675	97	65,353	3.0	6.0	Run body
220	36.6			62	86	5,313	3.0	4.0	Run tail
221	36.6			178	74	13,173	1.0	1.5	Riffle
222	36.6			181	71	12,919	3.0	5.0	Run head
223	36.4			1047	90	94,576	6.5	8.3	Run body
224	36.3			115	97	11,107	3.0	3.5	Run tail
225	36.3			224	92	20,644	1.5	2.0	Riffle
226	36.3			69	79	5,484	2.0	2.5	Run head
227	36.3			213	65	13,878	2.0	2.5	Run body
228	36.2			70	58	4,092	1.5	2.0	Run tail
229	36.2			74	54	4,022	1.2	2.0	Riffle
230	36.2			89	72	6,363	4.0	9.8	Pool head
231	36.2			175	131	22,846	6.0	12.3	Pool body
232	36.2			106	107	11,336	4.0	6.0	Pool tail
233	36.1			211	78	16,529	2.0	3.0	Pool head
234	35.7			2458	72	177,862	9.0	13.4	Pool body
235	35.6			210	53	11,010	3.0	3.5	Pool tail
236	35.5			353	97	34,136	1.0	1.5	Riffle
237	35.5			368	126	46,431	2.0	3.0	Run head
238	35.2			1394	100	139,804	3.5	7.0	Run body
239	35.2			48	84	4,006	3.0	4.0	Run tail
240	35.2			81	79	6,351	2.0	3.0	Riffle
241	35.2			70	60	4,157	3.0	4.0	Run head
242	35.2			74	68	5,054	4.5	5.8	Run body
243	35.1			62	65	3,996	1.5	2.0	Run tail
244	35.1			501	54	27,305	2.0	3.0	Riffle
245	35.0			79	82	6,466	1.5	2.5	Run head
246	35.0			302	65	19,636	2.0	3.0	Run body
247	35.0			114	31	3,548	1.5	2.0	Run tail
248	34.9			62	50	3,125	1.5	2.0	Riffle
249	34.9			151	50	7,602	3.0	4.0	Run head
250	34.7			1255	62	78,340	3.5	7.0	Run body
251	34.6			351	66	23,058	6.5	10.5	Pool body
252	34.6			119	82	9,791	3.0	4.0	Pool tail
253	34.5			293	77	22,628	1.0	2.0	Riffle
254	34.5			61	63	3,879	8.0	12.0	Pool head
255	34.4			445	79	35,344	4.0	8.0	Pool body
256	34.1			1722	91	157,333	3.0	4.0	Run body
257	34.1			137	81	11,136	1.5	2.0	Run tail
258	34.1			130	70	9,152	1.0	1.5	Riffle
259	34.0			103	79	8,137	2.0	2.5	Run head
260	34.0			452	59	26,907	2.5	3.5	Run body
261	33.9			142	38	5,468	1.5	2.0	Run tail

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft ²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
262	33.8			505	32	16,314	1.0	1.5	Riffle
263	33.8			86	53	4,509	2.0	2.5	Run head
264	33.8			265	52	13,757	3.0	3.5	Run body
265	33.8			59	57	3,342	2.0	2.5	Run tail
266	33.7			524	43	22,663	2.0	4.0	Riffle
267	33.6			241	67	16,237	3.0	4.0	Run head
268	33.5			690	116	79,804	2.5	5.0	Run body
269	33.4			231	79	18,336	1.0	2.0	Run tail
270	33.4			163	63	10,208	1.0	1.5	Riffle
271	33.4			49	74	3,588	6.0	7.5	Pool head
272	33.2			898	71	63,477	9.0	12.0	Pool body
273	33.2			102	39	3,988	2.0	3.0	Pool tail
274	33.2			190	55	10,514	1.0	1.5	Riffle
275	33.2			103	71	7,311	1.5	2.5	Run head
276	33.1			343	105	35,908	2.0	2.5	Run body
277	33.1			136	118	16,054	1.5	2.0	Run tail
278	33.0			312	62	19,368	1.0	1.5	Riffle
279	33.0			209	35	7,298	3.5	6.0	Run head
280	32.1			4454	174	776,561	5.5	9.2	Run body
281	32.1			143	124	17,763	4.0	5.5	Run tail
282	32.0			293	100	29,228	1.0	1.5	Riffle
283	32.0			163	107	17,489	2.5	3.0	Run head
284	32.0			294	86	25,244	3.5	4.0	Run body
285	31.9			41	86	3,565	2.0	3.7	Run tail
286	31.9			290	87	25,317	1.0	2.0	Riffle
287	31.9			157	43	6,710	2.5	3.0	Run head
288	31.7			838	55	45,952	3.5	5.0	Run body
289	31.7			112	85	9,543	2.5	3.0	Run tail
290	31.6			181	100	18,051	1.0	2.0	Riffle
291	31.6			148	108	15,990	4.0	5.5	Run head
292	31.5			475	89	42,320	5.0	6.0	Run body
293	31.5			154	62	9,597	1.5	2.5	Run tail
294	31.5			175	74	13,012	1.0	1.5	Riffle
295	31.4			210	100	21,058	3.0	4.5	Run head
296	31.3			567	87	49,612	4.0	5.5	Run body
297	31.3			139	54	7,465	2.5	4.0	Run tail
298	31.2			538	44	23,863	1.5	2.5	Riffle
299	31.2			122	70	8,583	3.5	4.5	Run head
300	31.1			240	61	14,568	3.5	5.0	Run body
301	31.1			41	72	2,974	2.0	3.0	Run tail
302	31.1			206	66	13,664	1.3	2.0	Riffle
303	31.1			98	75	7,324	3.0	4.0	Run head
304	30.7			1892	85	160,847	4.0	5.5	Run body
305	30.7			200	102	20,508	1.5	2.5	Run tail

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

Sampling Unit	RM	March 2010 BCE site	August 2010 BCE site	Length (ft)	Average width (ft)	Area (ft²)	Average depth (ft)	Maximum depth (ft)	July 2008 habitat type
306	30.6			113	83	9,452	1.2	2.0	Riffle
307	30.6			113	69	7,775	2.0	3.5	Run head
308	30.5			513	74	37,874	3.5	6.5	Run body
309	30.5			157	95	14,947	2.5	3.5	Run tail
310	30.4			259	37	9,478	1.0	2.0	Riffle
311	30.4			71	40	2,836	2.5	3.0	Run head
312	30.4			188	47	8,790	2.5	3.0	Run body
313	30.4			59	49	2,887	1.5	3.0	Run tail
314	30.2			946	43	40,519	1.2	2.0	Riffle
315	30.2			263	49	12,952	2.5	3.0	Run head
316	30.1			123	60	7,371	2.5	5.0	Run body
317	30.1			52	71	3,674	2.0	3.0	Run tail
318	30.1			189	298	56,219	1.5	2.0	Riffle
319	30.0			329	171	56,219	2.0	3.0	Run head
320	29.7			1444	155	224,395	5.0	8.0	Run body
321	29.7			68	59	3,978	3.0	4.0	Run tail
322	29.6			681	329	223,763	11.0	15.7	Pool body
323	29.6			222	84	18,626	3.0	7.0	Pool tail
324	29.5			109	38	4,188	1.0	2.0	Riffle
325	29.5			110	55	6,041	4.0	5.0	Run head
326	29.5			190	51	9,726	3.0	4.0	Run body
327	29.5			52	63	3,270	2.0	3.0	Run tail
328	29.5			70	58	4,066	1.2	2.0	Riffle
329	29.4			88	40	3,575	3.5	4.0	Run head
330	29.4			301	53	15,958	3.5	4.5	Run body
331	29.4			169	79	13,387	1.5	2.5	Run tail
332	29.3			192	168	32,257	1.2	2.0	Riffle
333	29.3			131	139	18,145	2.0	3.8	Run head
334	29.2			402	110	44,240	3.0	5.0	Run body
335	29.2			51	135	6,896	2.0	3.5	Run tail
336	29.2			247	92	22,792	1.0	1.5	Riffle
337	29.1			103	88	9,057	2.5	3.0	Run head
338	29.1			168	89	14,954	3.5	4.5	Run body
339	29.0			331	127	42,219	2.0	2.5	Run tail
340	29.0			447	90	40,119	1.5	2.0	Riffle

Table D-2. Percent cover and type for habitat units within the study area.

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
51.8	1	Pool head	7/8/2008	90	5			5	
51.7	2	Pool body	7/8/2008	80					20
51.7	3	Pool tail	7/8/2008	100					
51.6	4	Pool head	7/8/2008	100					
51.6	5	Pool body	7/8/2008	90					10
51.5	6	Pool tail	7/8/2008	100					
51.5	7	Riffle	7/8/2008	90	5			5	
51.4	8	Run head	7/8/2008	85				5	10
51.1	9	Run body	7/8/2008	60	10				30
51.0	10	Run tail	7/8/2008	90					10
50.9	11	Pool body	7/8/2008	50					50
50.8	12	Run body	7/8/2008	45	5				50
50.8	13	Run tail	7/8/2008	90				10	
50.6	14	Riffle	7/8/2008	80	10		10		
50.6	15	Run head	7/8/2008	90	10				
50.5	16	Run body	7/8/2008	95				5	
50.4	17	Run tail	7/8/2008	90				5	
50.3	18	Riffle	7/8/2008	90	5				5
50.3	19	Run head	7/8/2008	90					10
50.1	20	Run body	7/8/2008	95				5	
50.1	21	Run tail	7/8/2008	90	5			5	
50.1	22	Riffle	7/8/2008	95					5
50.0	23	Run head	7/8/2008	95				5	
49.9	24	Run body	7/8/2008	95				5	
49.8	25	Run tail	7/8/2008	95				5	
49.7	26	Riffle	7/8/2008	90	5			5	
49.7	27	Pool head	7/8/2008	85	10			5	
49.6	28	Pool body	7/8/2008	85	10			5	
49.6	29	Pool tail	7/8/2008	85	10			5	
49.6	30	Run head	7/8/2008	100					
49.3	31	Run body	7/8/2008	95		5			
49.3	32	Run tail	7/8/2008	95				5	
49.2	33	Riffle	7/8/2008	90	5			5	
49.2	34	Run head	7/8/2008	85	5			10	
49.1	35	Run body	7/8/2008	85	5			10	
49.1	36	Run tail	7/8/2008	95				5	
49.1	37	Riffle	7/8/2008	95				5	
49.1	38	Run head	7/8/2008	90		5		5	
49.1	39	Run body	7/8/2008	90	5			5	
49.0	40	Run tail	7/8/2008	95				5	
48.8	41	Riffle	7/8/2008	95				5	
48.8	42	Run head	7/8/2008	75				5	20
48.7	43	Run body	7/8/2008	90				10	

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
48.7	44	Run tail	7/8/2008	95				5	
48.4	45	Riffle	7/8/2008	90				10	
48.4	46	Run head	7/8/2008	90				10	
48.3	47	Run body	7/8/2008	90				10	
48.2	48	Run tail	7/8/2008	90				10	
48.2	49	Riffle	7/8/2008	90				10	
48.2	50	Run head	7/8/2008	90		5		5	
48.1	51	Run body	7/8/2008	95	5				
48.1	52	Run tail	7/8/2008	95	5				
48.0	53	Riffle	7/8/2008	95				5	
48.0	54	Pool head	7/8/2008	85	10			5	
47.2	55	Pool body	7/8/2008	85	10			5	
47.2	56	Pool tail	7/8/2008	95				5	
47.1	57	Riffle	7/8/2008	100					
47.0	58	Run head	7/8/2008	100					
46.9	59	Run body	7/8/2008	95				5	
46.9	60	Run tail	7/8/2008	90				10	
46.9	61	Riffle	7/8/2008	95				5	
46.9	62	Run head	7/8/2008	90				10	
46.8	63	Run body	7/8/2008	95				5	
46.8	64	Run tail	7/8/2008	95				5	
46.8	65	Riffle	7/8/2008	95				5	
46.8	66	Run head	7/8/2008	100					
46.0	67	Run body	7/8/2008	95				5	
46.0	68	Run tail	7/8/2008	95				5	
45.9	69	Run body	7/8/2008	100					
45.9	70	Riffle	7/8/2008	90				10	
45.9	71	Run head	7/8/2008	95				5	
45.8	72	Run body	7/8/2008	95				5	
45.8	73	Run tail	7/8/2008	100					
45.7	74	Riffle	7/8/2008	95				5	
45.7	75	Run head	7/9/2008	90				10	
45.7	76	Run body	7/9/2008	90				10	
45.7	77	Run tail	7/9/2008	100					
45.6	78	Riffle	7/9/2008	95				5	
45.6	79	Run head	7/9/2008	85				5	10
45.4	80	Run body	7/9/2008	80	15			5	
45.3	81	Pool body	7/9/2008	40		5		5	50
45.3	82	Run head	7/9/2008	45				5	50
45.1	83	Run body	7/9/2008	35		5		10	50
45.1	84	Run tail	7/9/2008	75		5		20	
45.0	85	Riffle	7/9/2008	70		5		25	
45.0	86	Pool head	7/9/2008	85		5		10	
44.9	87	Pool body	7/9/2008	90		5		5	
44.9	88	Pool tail	7/9/2008	95					5

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
44.8	89	Riffle	7/9/2008	90				10	
44.8	90	Run head	7/9/2008	90		5		5	
44.8	91	Run body	7/9/2008	100					
44.8	92	Run tail	7/9/2008	85				15	
44.7	93	Riffle	7/9/2008	80				20	
44.7	94	Run head	7/9/2008	90				10	
44.7	95	Run body	7/9/2008	100					
44.7	96	Run tail	7/9/2008	95				5	
44.6	97	Riffle	7/9/2008	90				10	
44.6	98	Run head	7/9/2008	95				5	
44.6	99	Run body	7/9/2008	95				5	
44.5	100	Run tail	7/9/2008	95				5	
44.5	101	Riffle	7/9/2008	95				5	
44.5	102	Run head	7/9/2008	100					
43.9	103	Run body	7/9/2008	90				10	
43.7	104	Pool body	7/9/2008	65				5	30
43.3	105	Run body	7/9/2008	65				5	30
43.3	106	Run tail	7/9/2008	90				5	5
43.2	107	Riffle	7/9/2008	85		5		10	
43.2	108	Run head	7/9/2008	95				5	
43.1	109	Run body	7/9/2008	95				5	
43.1	110	Run tail	7/9/2008	90				10	
43.0	111	Riffle	7/9/2008	95				5	
43.0	112	Pool head	7/9/2008	65		5			30
43.0	113	Pool body	7/9/2008	60		10			30
43.0	114	Pool tail	7/9/2008	70		25		5	
43.0	115	Run head	7/9/2008	70		20		10	
42.9	116	Run body	7/9/2008	100					
42.9	117	Run tail	7/9/2008	95				5	
42.9	118	Riffle	7/9/2008	95				5	
42.9	119	Run head	7/9/2008	95				5	
42.7	120	Run body	7/9/2008	95				5	
42.7	121	Run tail	7/9/2008	95				5	
42.7	122	Riffle	7/9/2008	90				5	5
42.7	123	Run head	7/9/2008	95				5	
42.4	124	Run body	7/9/2008	95				5	
42.4	125	Run body	7/9/2008	95				5	
42.3	126	Riffle	7/9/2008	80				20	
42.3	127	Run body	7/9/2008	100					
42.3	128	Riffle	7/9/2008	75	5	5		15	
42.2	129	Run head	7/9/2008	90				10	
42.1	130	Run body	7/9/2008	90				10	
42.0	131	Run tail	7/9/2008	95				5	
41.9	132	Riffle	7/9/2008	95				5	
41.9	133	Run head	7/9/2008	95				5	

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
41.8	134	Run body	7/9/2008	95				5	
41.8	135	Run tail	7/9/2008	95				5	
41.7	136	Riffle	7/9/2008	95				5	
41.7	137	Run head	7/9/2008	90				10	
41.2	138	Run body	7/9/2008	100					
41.2	139	Run tail	7/9/2008	95				5	
41.1	140	Riffle	7/9/2008	95				5	
41.1	141	Run head	7/9/2008	80					20
41.0	142	Run body	7/9/2008	95				5	
41.0	143	Run tail	7/9/2008	95				5	
40.9	144	Riffle	7/9/2008	95				5	
40.9	145	Run head	7/9/2008	100					
40.5	146	Run body	7/9/2008	65				10	25
40.5	147	Run tail	7/9/2008	85				15	
40.4	148	Riffle	7/9/2008	70				30	
40.4	149	Run head	7/9/2008	75				5	20
40.3	150	Run body	7/9/2008	100					
40.3	151	Run tail	7/9/2008	100					
40.2	152	Riffle	7/9/2008	95				5	
40.2	153	Run head	7/9/2008	100					
39.7	154	Run body	7/9/2008	95				5	
39.7	155	Run tail	7/9/2008	95				5	
39.7	156	Riffle	2/10/2009	95					5
39.6	157	Run head	2/10/2009	100					
39.5	158	Run body	2/10/2009	80					20
39.5	159	Run tail	2/10/2009	80					20
39.4	160	Riffle	2/10/2009	95					5
39.4	161	Run head	2/10/2009	95					
39.3	162	Run body	2/10/2009	95				5	
39.3	163	Run tail	2/10/2009	95				5	
39.2	164	Riffle	2/10/2009	95					5
39.2	165	Pool head	2/10/2009	100					
38.9	166	Pool body	2/10/2009	90					10
38.9	167	Pool tail	2/10/2009	100					
38.9	168	Riffle	2/10/2009	100					
38.9	169	Run head	2/10/2009	100					
38.8	170	Run body	2/10/2009	100					
38.8	171	Pool body	2/10/2009	90				5	5
38.8	172	Run head	2/10/2009	95				5	
38.7	173	Run body	2/10/2009	95				5	
38.7	174	Run tail	2/10/2009	100					
38.7	175	Riffle	2/10/2009	100					
38.6	176	Run head	2/10/2009	100					
38.6	177	Run body	2/10/2009	100					
38.6	178	Run tail	2/10/2009	100					

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
38.5	179	Riffle	2/10/2009	100					
38.5	180	Run head	2/10/2009	90					10
38.4	181	Run body	2/10/2009	100					
38.3	182	Pool body	2/10/2009	80					20
38.3	183	Pool tail	2/10/2009	90				5	5
38.3	184	Run head	2/10/2009	100					
38.2	185	Run body	2/10/2009	100					
38.2	186	Run tail	2/10/2009	100					
38.2	187	Riffle	2/10/2009	95				5	
38.1	188	Pool head	2/10/2009	95				5	
38.1	189	Pool body	2/11/2009	90					10
38.1	190	Pool tail	2/11/2009	100					
38.1	191	Riffle	2/11/2009	100					
38.1	192	Pool head	2/11/2009	90					10
38.0	193	Pool body	2/11/2009	70					30
38.0	194	Pool tail	2/11/2009	100					
38.0	195	Run head	2/11/2009	100					
37.9	196	Run body	2/11/2009	100					
37.9	197	Run tail	2/11/2009	100					
37.8	198	Riffle	2/11/2009	100					
37.8	199	Pool head	2/11/2009	85		15			
37.7	200	Pool body	2/11/2009	100					
37.6	201	Pool tail	2/11/2009	100					
37.6	202	Riffle	2/11/2009	100					
37.6	203	Run head	2/11/2009	100					
37.5	204	Run body	2/11/2009	100					
37.4	205	Run tail	2/11/2009	100					
37.3	206	Riffle	2/11/2009	100					
37.3	207	Run head	2/11/2009	100					
37.1	208	Run body	2/11/2009	100					
37.1	209	Run tail	2/11/2009	100					
37.0	210	Riffle	2/11/2009	100					
37.0	211	Run head	2/11/2009	100					
36.9	212	Run body	2/11/2009	100					
36.9	213	Run tail	2/11/2009	100					
36.9	214	Pool head	2/11/2009	100					
36.9	215	Pool body	2/11/2009	100					
36.9	216	Pool tail	2/11/2009	100					
36.8	217	Riffle	2/11/2009	100					
36.8	218	Run head	2/11/2009	100					
36.6	219	Run body	2/11/2009	100					
36.6	220	Run tail	2/11/2009	100					
36.6	221	Riffle	2/11/2009	100					
36.6	222	Run head	2/11/2009	100					
36.4	223	Run body	2/11/2009	100					

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
36.3	224	Run tail	2/11/2009	100					
36.3	225	Riffle	2/11/2009	100					
36.3	226	Run head	2/11/2009	100					
36.3	227	Run body	2/11/2009	100					
36.2	228	Run tail	2/11/2009	100					
36.2	229	Riffle	2/11/2009	100					
36.2	230	Pool head	2/11/2009	100					
36.2	231	Pool body	2/11/2009	100					
36.2	232	Pool tail	2/11/2009	100					
36.1	233	Pool head	2/11/2009	100					
35.7	234	Pool body	2/11/2009	100					
35.6	235	Pool tail	2/11/2009	100					
35.5	236	Riffle	2/11/2009	100					
35.5	237	Run head	2/11/2009	100					
35.2	238	Run body	2/11/2009	100					
35.2	239	Run tail	2/12/2009	95				5	
35.2	240	Riffle	2/12/2009	100					
35.2	241	Run head	2/12/2009	100					
35.2	242	Run body	2/12/2009	100					
35.1	243	Run tail	2/12/2009	100					
35.1	244	Riffle	2/12/2009	100					
35.0	245	Run head	2/12/2009	95				5	
35.0	246	Run body	2/12/2009	95				5	
35.0	247	Run tail	2/12/2009	100					
34.9	248	Riffle	2/12/2009	100					
34.9	249	Run head	2/12/2009	95		5			
34.7	250	Run body	2/12/2009	100					
34.6	251	Pool body	2/12/2009	75				5	20
34.6	252	Pool tail	2/12/2009	100					
34.5	253	Riffle	2/12/2009	95				5	
34.5	254	Pool head	2/12/2009	100					
34.4	255	Pool body	2/12/2009	100					
34.1	256	Run body	2/12/2009	100					
34.1	257	Run tail	2/12/2009	95				5	
34.1	258	Riffle	2/12/2009	100					
34.0	259	Run head	2/12/2009	100					
34.0	260	Run body	2/12/2009	100					
33.9	261	Run tail	2/12/2009	100					
33.8	262	Riffle	2/12/2009	100					
33.8	263	Run head	2/12/2009	100					
33.8	264	Run body	2/12/2009	100					
33.8	265	Run tail	2/12/2009	100					
33.7	266	Riffle	2/12/2009	100					
33.6	267	Run head	2/12/2009	100					
33.5	268	Run body	2/12/2009	100					

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
33.4	269	Run tail	2/12/2009	100					
33.4	270	Riffle	2/12/2009	100					
33.4	271	Pool head	2/12/2009	100					
33.2	272	Pool body	2/12/2009	70					30
33.2	273	Pool tail	2/12/2009	100					
33.2	274	Riffle	2/12/2009	100					
33.2	275	Run head	2/12/2009	100					
33.1	276	Run body	2/12/2009	95					5
33.1	277	Run tail	2/12/2009	100					
33.0	278	Riffle	2/12/2009	100					
33.0	279	Run head	2/12/2009	100					
32.1	280	Run body	2/12/2009	60					40
32.1	281	Run tail	2/12/2009						
32.0	282	Riffle	2/12/2009						
32.0	283	Run head	2/12/2009						
32.0	284	Run body	2/12/2009						
31.9	285	Run tail	2/12/2009						
31.9	286	Riffle	2/12/2009						
31.9	287	Run head	2/12/2009						
31.7	288	Run body	2/12/2009						
31.7	289	Run tail	2/12/2009						
31.6	290	Riffle	2/12/2009						
31.6	291	Run head	2/12/2009						
31.5	292	Run body	2/12/2009						
31.5	293	Run tail	2/12/2009						
31.5	294	Riffle	2/12/2009	100					
31.4	295	Run head	2/12/2009	100					
31.3	296	Run body	2/12/2009	100					
31.3	297	Run tail	2/12/2009	100					
31.2	298	Riffle	2/12/2009	100					
31.2	299	Run head	2/13/2009	100					
31.1	300	Run body	2/13/2009	100					
31.1	301	Run tail	2/13/2009	100					
31.1	302	Riffle	2/13/2009	100					
31.1	303	Run head	2/13/2009	100					
30.7	304	Run body	2/13/2009	100					
30.7	305	Run tail	2/13/2009	90					10
30.6	306	Riffle	2/13/2009	100					
30.6	307	Run head	2/13/2009	100					
30.5	308	Run body	2/13/2009	100					
30.5	309	Run tail	2/13/2009	100					
30.4	310	Riffle	2/13/2009	85				15	
30.4	311	Run head	2/13/2009	100					
30.4	312	Run body	2/13/2009	100					
30.4	313	Run tail	2/13/2009	100					

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Sampling unit	Habitat type	Habitat survey date	No cover (%)	Boulder (%)	Wood (%)	Ledge (%)	Overhang (%)	Aquatic vegetation (%)
30.2	314	Riffle	2/13/2009	90				10	
30.2	315	Run head	2/13/2009	100					
30.1	316	Run body	2/13/2009	100					
30.1	317	Run tail	2/13/2009	100					
30.1	318	Riffle	2/13/2009	100					
30.0	319	Run head	2/13/2009	100					
29.7	320	Run body	2/13/2009	70					30
29.7	321	Run tail	2/13/2009	90					10
29.6	322	Pool body	2/13/2009	100					
29.6	323	Pool tail	2/13/2009	100					
29.5	324	Riffle	2/13/2009	100					
29.5	325	Run head	2/13/2009	95	5				
29.5	326	Run body	2/13/2009	85					15
29.5	327	Run tail	2/13/2009	100					
29.5	328	Riffle	2/13/2009	100					
29.4	329	Run head	2/13/2009	100					
29.4	330	Run body	2/13/2009	100					
29.4	331	Run tail	2/13/2009	100					
29.3	332	Riffle	2/13/2009	90				10	
29.3	333	Run head	2/13/2009	100					
29.2	334	Run body	2/13/2009	100					
29.2	335	Run tail	2/13/2009	100					
29.2	336	Riffle	2/13/2009	100					
29.1	337	Run head	2/13/2009	100					
29.1	338	Run body	2/13/2009	90					10
29.0	339	Run tail	2/13/2009	100					
29.0	340	Riffle	2/13/2009	100					

Table D-3. Substrate types for sampling units within the study area.

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
51.8	1	Pool head	7/8/2008	10	50	40				
51.7	2	Pool body	7/8/2008	50	40	10				
51.7	3	Pool tail	7/8/2008	20	30	50				
51.6	4	Pool head	7/8/2008	50	20	30				
51.6	5	Pool body	7/8/2008	50	20	25		5		
51.5	6	Pool tail	7/8/2008	40	30	30				
51.5	7	Riffle	7/8/2008		30	60	10			
51.4	8	Run head	7/8/2008		20	60	10	10		
51.1	9	Run body	7/8/2008	15	15	60	10			
51.0	10	Run tail	7/8/2008			60	30	10		
50.9	11	Pool body	7/8/2008	20	10	50		20		
50.8	12	Run body	7/8/2008	20	10	50		20		
50.8	13	Run tail	7/8/2008			60	30	10		
50.6	14	Riffle	7/8/2008			60	30	10		
50.6	15	Run head	7/8/2008		10	50	40			
50.5	16	Run body	7/8/2008	10	10	60	20			
50.4	17	Run tail	7/8/2008		20	60	20			
50.3	18	Riffle	7/8/2008		20	60	20			
50.3	19	Run head	7/8/2008		20	60	20			
50.1	20	Run body	7/8/2008		20	60	20			
50.1	21	Run tail	7/8/2008		20	60	20			
50.1	22	Riffle	7/8/2008		20	60	20			
50.0	23	Run head	7/8/2008		20	60	20			
49.9	24	Run body	7/8/2008		60	20	20			
49.8	25	Run tail	7/8/2008		40	40	20			
49.7	26	Riffle	7/8/2008		20	60	20			
49.7	27	Pool head	7/8/2008	20	20	40	10	10		
49.6	28	Pool body	7/8/2008	20	20	40	10	10		
49.6	29	Pool tail	7/8/2008	10	20	60	10			
49.6	30	Run head	7/8/2008		20	60	20			
49.3	31	Run body	7/8/2008		20	60	20			
49.3	32	Run tail	7/8/2008		10	70	20			
49.2	33	Riffle	7/8/2008		10	70	20			
49.2	34	Run head	7/8/2008		10	70	20			
49.1	35	Run body	7/8/2008		10	70	20			
49.1	36	Run tail	7/8/2008		10	70	20			
49.1	37	Riffle	7/8/2008		10	70	20			
49.1	38	Run head	7/8/2008		10	70	20			
49.1	39	Run body	7/8/2008		10	70	20			
49.0	40	Run tail	7/8/2008		10	70	20			
48.8	41	Riffle	7/8/2008		10	70	20			
48.8	42	Run head	7/8/2008		10	70	20			
48.7	43	Run body	7/8/2008		40	40	20			

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
48.7	44	Run tail	7/8/2008		40	40	20			
48.4	45	Riffle	7/8/2008		20	60	20			
48.4	46	Run head	7/8/2008		10	40	50			
48.3	47	Run body	7/8/2008		10	50	40			
48.2	48	Run tail	7/8/2008		10	70	20			
48.2	49	Riffle	7/8/2008		10	70	20			
48.2	50	Run head	7/8/2008		10	70	20			
48.1	51	Run body	7/8/2008	20	10	50	20			
48.1	52	Run tail	7/8/2008	20	10	50	20			
48.0	53	Riffle	7/8/2008		10	70	20			
48.0	54	Pool head	7/8/2008	20	10	60	5	5		
47.2	55	Pool body	7/8/2008	20	10	60	5	5		
47.2	56	Pool tail	7/8/2008		10	70	20			
47.1	57	Riffle	7/8/2008		10	70	20			
47.0	58	Run head	7/8/2008		10	70	20			
46.9	59	Run body	7/8/2008	20	10	50	20			
46.9	60	Run tail	7/8/2008		20	60	20			
46.9	61	Riffle	7/8/2008		10	70	20			
46.9	62	Run head	7/8/2008		10	70	20			
46.8	63	Run body	7/8/2008		10	70	20			
46.8	64	Run tail	7/8/2008		10	60	30			
46.8	65	Riffle	7/8/2008		10	60	30			
46.8	66	Run head	7/8/2008		10	50	30	10		
46.0	67	Run body	7/8/2008		20	50	20	10		
46.0	68	Run tail	7/8/2008		10	70	20			
45.9	69	Run body	7/8/2008		10	70	20			
45.9	70	Riffle	7/8/2008			20	70	10		
45.9	71	Run head	7/8/2008			30	40	30		
45.8	72	Run body	7/8/2008			40	40	20		
45.8	73	Run tail	7/8/2008			40	50	10		
45.7	74	Riffle	7/8/2008			40	50	10		
45.7	75	Run head	7/9/2008		10	60	20	10		
45.7	76	Run body	7/9/2008		10	60	20	10		
45.7	77	Run tail	7/9/2008		10	60	20	10		
45.6	78	Riffle	7/9/2008			70	20	10		
45.6	79	Run head	7/9/2008		10	10	30	50		
45.4	80	Run body	7/9/2008	20	20	30		30		
45.3	81	Pool body	7/9/2008	30	20	20		30		
45.3	82	Run head	7/9/2008			10	30	50	10	
45.1	83	Run body	7/9/2008	10	20	50	10	10		
45.1	84	Run tail	7/9/2008		10	70	20			
45.0	85	Riffle	7/9/2008		10	60	30			
45.0	86	Pool head	7/9/2008		10	60	30			
44.9	87	Pool body	7/9/2008			60	20	20		
44.9	88	Pool tail	7/9/2008			60	20	20		

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
44.8	89	Riffle	7/9/2008		20	60	20			
44.8	90	Run head	7/9/2008			40	50	10		
44.8	91	Run body	7/9/2008		10	60	30			
44.8	92	Run tail	7/9/2008		10	60	30			
44.7	93	Riffle	7/9/2008			60	30	10		
44.7	94	Run head	7/9/2008			60	30	10		
44.7	95	Run body	7/9/2008							
44.7	96	Run tail	7/9/2008			40	10	50		
44.6	97	Riffle	7/9/2008		10	50	40			
44.6	98	Run head	7/9/2008		10	50	40			
44.6	99	Run body	7/9/2008		10	40	40	10		
44.5	100	Run tail	7/9/2008		10	40	40	10		
44.5	101	Riffle	7/9/2008	10	10	50	30			
44.5	102	Run head	7/9/2008		10	50	30	10		
43.9	103	Run body	7/9/2008	40	10	30	10	10		
43.7	104	Pool body	7/9/2008	20	10	20		50		
43.3	105	Run body	7/9/2008	20	10	20		50		
43.3	106	Run tail	7/9/2008		10	60	20	10		
43.2	107	Riffle	7/9/2008		10	60	30			
43.2	108	Run head	7/9/2008		10	60	20	10		
43.1	109	Run body	7/9/2008		10	60	30			
43.1	110	Run tail	7/9/2008		10	60	30			
43.0	111	Riffle	7/9/2008		10	60	30			
43.0	112	Pool head	7/9/2008		10	50	30	10		
43.0	113	Pool body	7/9/2008		10	50	30	10		
43.0	114	Pool tail	7/9/2008		10	50	30	10		
43.0	115	Run head	7/9/2008		10	50	30	10		
42.9	116	Run body	7/9/2008		10	60	30			
42.9	117	Run tail	7/9/2008		10	60	30			
42.9	118	Riffle	7/9/2008		10	60	30			
42.9	119	Run head	7/9/2008		20	50	30			
42.7	120	Run body	7/9/2008		20	50	30			
42.7	121	Run tail	7/9/2008		10	60	30			
42.7	122	Riffle	7/9/2008		10	50	40			
42.7	123	Run head	7/9/2008		10	50	40			
42.4	124	Run body	7/9/2008		10	50	40			
42.4	125	Run body	7/9/2008		10	50	40			
42.3	126	Riffle	7/9/2008		10	50	40			
42.3	127	Run body	7/9/2008	50		40	10			
42.3	128	Riffle	7/9/2008	15	10	50	20	5		
42.2	129	Run head	7/9/2008	15	10	50	20	5		
42.1	130	Run body	7/9/2008		10	60	30			
42.0	131	Run tail	7/9/2008		10	50	40			
41.9	132	Riffle	7/9/2008		15	50	35			
41.9	133	Run head	7/9/2008	15	15	45	25			

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
41.8	134	Run body	7/9/2008	15	15	40	20	10		
41.8	135	Run tail	7/9/2008		10	60	30			
41.7	136	Riffle	7/9/2008		10	60	30			
41.7	137	Run head	7/9/2008	15	10	50	25			
41.2	138	Run body	7/9/2008	15	10	50	25			
41.2	139	Run tail	7/9/2008		10	60	20	10		
41.1	140	Riffle	7/9/2008		10	50	30	10		
41.1	141	Run head	7/9/2008		10	50	30	10		
41.0	142	Run body	7/9/2008		10	50	30	10		
41.0	143	Run tail	7/9/2008		10	60	20	10		
40.9	144	Riffle	7/9/2008		10	60	20	10		
40.9	145	Run head	7/9/2008		10	50	40			
40.5	146	Run body	7/9/2008		50	20		30		
40.5	147	Run tail	7/9/2008		10	60	30			
40.4	148	Riffle	7/9/2008		10	50	40			
40.4	149	Run head	7/9/2008		10	50	30	10		
40.3	150	Run body	7/9/2008							
40.3	151	Run tail	7/9/2008		20	50	30			
40.2	152	Riffle	7/9/2008		20	50	30			
40.2	153	Run head	7/9/2008		20	50	30			
39.7	154	Run body	7/9/2008	20	10	50	10	10		
39.7	155	Run tail	7/9/2008		10	50	40			
39.7	156	Riffle	2/10/2009			50	40	10		
39.6	157	Run head	2/10/2009			30	20	50		
39.5	158	Run body	2/10/2009			30	20	50		
39.5	159	Run tail	2/10/2009			30	20	50		
39.4	160	Riffle	2/10/2009			50	40	10		
39.4	161	Run head	2/10/2009		10	50	30	10		
39.3	162	Run body	2/10/2009		10	50	30	10		
39.3	163	Run tail	2/10/2009	5		55	30	10		
39.2	164	Riffle	2/10/2009			50	40	10		
39.2	165	Pool head	2/10/2009			30	60	10		
38.9	166	Pool body	2/10/2009			20	50	30		
38.9	167	Pool tail	2/10/2009			50	40	10		
38.9	168	Riffle	2/10/2009			50	40	10		
38.9	169	Run head	2/10/2009			60	25	15		
38.8	170	Run body	2/10/2009			30	40	30		
38.8	171	Pool body	2/10/2009		5	60	20	15		
38.8	172	Run head	2/10/2009			60	30	10		
38.7	173	Run body	2/10/2009			60	30	10		
38.7	174	Run tail	2/10/2009			60	30	10		
38.7	175	Riffle	2/10/2009			60	30	10		
38.6	176	Run head	2/10/2009			60	30	10		
38.6	177	Run body	2/10/2009			60	30	10		
38.6	178	Run tail	2/10/2009			60	30	10		

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
38.5	179	Riffle	2/10/2009			60	30	10		
38.5	180	Run head	2/10/2009			50	20	30		
38.4	181	Run body	2/10/2009			60	30	10		
38.3	182	Pool body	2/10/2009		5	45	20	30		
38.3	183	Pool tail	2/10/2009		5	60	20	15		
38.3	184	Run head	2/10/2009			60	30	10		
38.2	185	Run body	2/10/2009			70	20	10		
38.2	186	Run tail	2/10/2009			60	30	10		
38.2	187	Riffle	2/10/2009			70	20	10		
38.1	188	Pool head	2/10/2009			60	30	10		
38.1	189	Pool body	2/11/2009		5	60	25	10		
38.1	190	Pool tail	2/11/2009			60	20	10	10	
38.1	191	Riffle	2/11/2009			70	20	10		
38.1	192	Pool head	2/11/2009			50	20	20	10	
38.0	193	Pool body	2/11/2009	20		20	30	30		
38.0	194	Pool tail	2/11/2009			40	40	20		
38.0	195	Run head	2/11/2009			50	40	10		
37.9	196	Run body	2/11/2009			60	30	10		
37.9	197	Run tail	2/11/2009			60	30	5	5	
37.8	198	Riffle	2/11/2009			60	30	10		
37.8	199	Pool head	2/11/2009			60	30	10		
37.7	200	Pool body	2/11/2009	10			60	30		
37.6	201	Pool tail	2/11/2009			5	75	20		
37.6	202	Riffle	2/11/2009	5		5	80	10		
37.6	203	Run head	2/11/2009			10	60	20	10	
37.5	204	Run body	2/11/2009			30	60	10		
37.4	205	Run tail	2/11/2009			40	60			
37.3	206	Riffle	2/11/2009			40	60			
37.3	207	Run head	2/11/2009			50	40	10		
37.1	208	Run body	2/11/2009			50	40	10		
37.1	209	Run tail	2/11/2009			50	50			
37.0	210	Riffle	2/11/2009			60	40			
37.0	211	Run head	2/11/2009			50	40	10		
36.9	212	Run body	2/11/2009			10	60	30		
36.9	213	Run tail	2/11/2009			20	70	10		
36.9	214	Pool head	2/11/2009			20	70	10		
36.9	215	Pool body	2/11/2009			20	50	30		
36.9	216	Pool tail	2/11/2009			10	60	30		
36.8	217	Riffle	2/11/2009			30	60	10		
36.8	218	Run head	2/11/2009			40	50	10		
36.6	219	Run body	2/11/2009			20	40	40		
36.6	220	Run tail	2/11/2009			20	60	20		
36.6	221	Riffle	2/11/2009			30	60	10		
36.6	222	Run head	2/11/2009			40	60			
36.4	223	Run body	2/11/2009			20	60	20		

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
36.3	224	Run tail	2/11/2009			30	60	10		
36.3	225	Riffle	2/11/2009			30	60	10		
36.3	226	Run head	2/11/2009			30	60	10		
36.3	227	Run body	2/11/2009			30	60	10		
36.2	228	Run tail	2/11/2009			30	60	10		
36.2	229	Riffle	2/11/2009			30	60	10		
36.2	230	Pool head	2/11/2009			30	60	10		
36.2	231	Pool body	2/11/2009			30	60	10		
36.2	232	Pool tail	2/11/2009			20	60	20		
36.1	233	Pool head	2/11/2009				80	20		
35.7	234	Pool body	2/11/2009	25		20	40	15		
35.6	235	Pool tail	2/11/2009			30	60	10		
35.5	236	Riffle	2/11/2009			30	60	10		
35.5	237	Run head	2/11/2009			30	60	10		
35.2	238	Run body	2/11/2009		5	15	20	60		
35.2	239	Run tail	2/12/2009			30	60	5	5	
35.2	240	Riffle	2/12/2009			35	60	5		
35.2	241	Run head	2/12/2009			35	60	5		
35.2	242	Run body	2/12/2009			30	65	5		
35.1	243	Run tail	2/12/2009			20	80			
35.1	244	Riffle	2/12/2009			20	60	20		
35.0	245	Run head	2/12/2009			20	70	10		
35.0	246	Run body	2/12/2009			40	50	10		
35.0	247	Run tail	2/12/2009			20	70	10		
34.9	248	Riffle	2/12/2009			10	80	10		
34.9	249	Run head	2/12/2009			20	70	10		
34.7	250	Run body	2/12/2009	5		25	60	10		
34.6	251	Pool body	2/12/2009	40		20	20	20		
34.6	252	Pool tail	2/12/2009	30		30	20	20		
34.5	253	Riffle	2/12/2009	5		30	65			
34.5	254	Pool head	2/12/2009	40		10	20	30		
34.4	255	Pool body	2/12/2009			30	50	20		
34.1	256	Run body	2/12/2009			30	60	10		
34.1	257	Run tail	2/12/2009			40	60			
34.1	258	Riffle	2/12/2009			30	60	10		
34.0	259	Run head	2/12/2009			40	50	10		
34.0	260	Run body	2/12/2009			30	40	30		
33.9	261	Run tail	2/12/2009			30	50	20		
33.8	262	Riffle	2/12/2009			30	60	10		
33.8	263	Run head	2/12/2009			40	60			
33.8	264	Run body	2/12/2009			40	50	10		
33.8	265	Run tail	2/12/2009			40	60			
33.7	266	Riffle	2/12/2009			40	50	10		
33.6	267	Run head	2/12/2009			10	70	20		
33.5	268	Run body	2/12/2009			20	40	40		

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)		
33.4	269	Run tail	2/12/2009			20	50	30				
33.4	270	Riffle	2/12/2009			30	60	10				
33.4	271	Pool head	2/12/2009			40	40	20				
33.2	272	Pool body	2/12/2009	10		20	30	30	10			
33.2	273	Pool tail	2/12/2009			40	50	10				
33.2	274	Riffle	2/12/2009			40	50	10				
33.2	275	Run head	2/12/2009			50	40	10				
33.1	276	Run body	2/12/2009			25	60	5	10			
33.1	277	Run tail	2/12/2009			40	50	10				
33.0	278	Riffle	2/12/2009			20	70	10				
33.0	279	Run head	2/12/2009			20	40	40				
32.1	280	Run body	2/12/2009				50	50				
32.1	281	Run tail	2/12/2009	No data collected								
32.0	282	Riffle	2/12/2009	No data collected								
32.0	283	Run head	2/12/2009	No data collected								
32.0	284	Run body	2/12/2009	No data collected								
31.9	285	Run tail	2/12/2009	No data collected								
31.9	286	Riffle	2/12/2009	No data collected								
31.9	287	Run head	2/12/2009	No data collected								
31.7	288	Run body	2/12/2009	No data collected								
31.7	289	Run tail	2/12/2009	No data collected								
31.6	290	Riffle	2/12/2009	No data collected								
31.6	291	Run head	2/12/2009	No data collected								
31.5	292	Run body	2/12/2009	No data collected								
31.5	293	Run tail	2/12/2009	No data collected								
31.5	294	Riffle	2/12/2009			40	50		10			
31.4	295	Run head	2/12/2009			20	70	10				
31.3	296	Run body	2/12/2009			10	60	30				
31.3	297	Run tail	2/12/2009			10	60	30				
31.2	298	Riffle	2/12/2009			30	60	10				
31.2	299	Run head	2/13/2009			40	50	10				
31.1	300	Run body	2/13/2009			30	40	30				
31.1	301	Run tail	2/13/2009			30	60	10				
31.1	302	Riffle	2/13/2009			30	60	10				
31.1	303	Run head	2/13/2009	10		40	40	10				
30.7	304	Run body	2/13/2009	10		40	40	10				
30.7	305	Run tail	2/13/2009			40	40	20				
30.6	306	Riffle	2/13/2009			40	50	10				
30.6	307	Run head	2/13/2009			40	50	10				
30.5	308	Run body	2/13/2009			40	50	10				
30.5	309	Run tail	2/13/2009			40	50	10				
30.4	310	Riffle	2/13/2009			30	50	20				
30.4	311	Run head	2/13/2009			30	60	10				
30.4	312	Run body	2/13/2009			40	50	10				
30.4	313	Run tail	2/13/2009		5	35	50	10				

*Population size estimates of O. mykiss
in the Lower Tuolumne River*

River mile	Unit	Habitat type	Habitat survey date	Bedrock (%)	Boulder (%)	Cobble (%)	Gravel (%)	Sand (%)	Silt (%)	Organic (%)
30.2	314	Riffle	2/13/2009			30	60	10		
30.2	315	Run head	2/13/2009			30	60	10		
30.1	316	Run body	2/13/2009			30	60	10		
30.1	317	Run tail	2/13/2009			30	60	10		
30.1	318	Riffle	2/13/2009			40	50	10		
30.0	319	Run head	2/13/2009			5	15	80		
29.7	320	Run body	2/13/2009				30	70		
29.7	321	Run tail	2/13/2009				30	70		
29.6	322	Pool body	2/13/2009				20	80		
29.6	323	Pool tail	2/13/2009				30	70		
29.5	324	Riffle	2/13/2009			30	60	10		
29.5	325	Run head	2/13/2009			40	60			
29.5	326	Run body	2/13/2009				20	80		
29.5	327	Run tail	2/13/2009				60	40		
29.5	328	Riffle	2/13/2009			30	70			
29.4	329	Run head	2/13/2009			20	60	10	10	
29.4	330	Run body	2/13/2009			10	70	20		
29.4	331	Run tail	2/13/2009			10	70	20		
29.3	332	Riffle	2/13/2009			10	80	10		
29.3	333	Run head	2/13/2009			10	70	20		
29.2	334	Run body	2/13/2009			20	70	10		
29.2	335	Run tail	2/13/2009			10	70	20		
29.2	336	Riffle	2/13/2009			10	80	10		
29.1	337	Run head	2/13/2009			10	60	30		
29.1	338	Run body	2/13/2009	15		30	30	25		
29.0	339	Run tail	2/13/2009	40		20	20	20		
29.0	340	Riffle	2/13/2009	20		10	60	10		

Appendix E: Water Quality Data

Table E-1. Water quality data for the sampling units selected for snorkel sampling, March 2010.

RM	Unit	Habitat type	Sample date	Start time	Water temperature (C)	DO (ppm)	Specific conductivity (mS)	Horizontal visability (ft)	Vertical visability (ft)	Average depth (ft)	Maximum depth (ft)
51.6	4	Pool Head	4-Mar	12:18	10.5	9.93	32.1	8.5		3.0	6.5
51.6	5	Pool Body	1-Mar	11:36	10.6	10.58	29.1	13.5	19.0	15.0	30.0
50.9	11	Pool Body	1-Mar	15:51	11.3	12.35	30.5	13.5	18.0	12.0	25.0
50.8	12	Run Body	1-Mar	15:30	11.3	12.35	30.5	13.5		6.0	10.0
50.6	15	Run Head	4-Mar	14:35	11.5	11.12	33.3	8.0		3.5	6.0
50.5	16	Run Body	2-Mar	10:41	10.6	10.64	28.1	17.0		7.0	11.0
50.3	18	Riffle	5-Mar	12:53	11.3	11.16	30.6	10.5		2.0	5.0
50.3	19	Run Head	5-Mar	13:52	11.3	11.16	30.6	10.5		4.0	8.0
50.1	20	Run Body	5-Mar	13:15	11.3	11.16	30.6	10.5		5.0	12.0
50.1	22	Riffle	2-Mar	16:10	11.0	11.53	32.5	17.0		1.5	4.0
49.7	26	Riffle	4-Mar	15:42	11.8	11.36	35.7	8.5		1.5	3.0
49.7	27	Pool Head	3-Mar	10:43	10.2	9.92	29.3	15.0		3.0	4.0
49.6	28	Pool Body	3-Mar	9:55	10.2	9.92	29.3	15.0		8.0	15.0
48.8	42	Run Head	3-Mar	14:05	10.6	11.18	30.6	15.0		1.5	2.5
48.7	43	Run Body	3-Mar	13:20	10.6	11.18	30.6	15.0		2.5	4.0
48.0	54	Pool Head	3-Mar	12:01	10.5	10.95	31.1	15.0		4.0	7.5
45.9	70	Riffle	5-Mar	10:59	10.6	10.38	37.4	10.5		2.0	3.5
45.0	86	Pool Head	6-Mar	10:44	10.7	10.59	37.4	12.0		5.0	11.0
44.8	90	Run Head	6-Mar	11:31	10.7	10.59	37.4	12.0		0.8	2.0
44.7	93	Riffle	6-Mar	11:52	12.3	11.59	39.4	9.0		2.0	4.0
44.5	101	Riffle	6-Mar	13:32	12.3	11.59	39.4	9.0		2.0	6.5
43.7	104	Pool Body	6-Mar	14:52	12.1	11.92	39.8	8.5	10.0	7.0	12.0
43.0	111	Riffle	7-Mar	10:02	11.5	10.78	39.9	11.5		1.5	3.0
43.0	112	Pool Head	6-Mar	16:24	12.1	11.70	40.6	9.0		2.0	4.0
43.0	113	Pool Body	6-Mar	16:07	12.1	11.70	40.6	9.0	10.0	5.0	10.0
42.9	116	Run Body	7-Mar	10:57	11.5	10.78	39.9	11.5		5.0	10.0
42.9	119	Run Head	7-Mar	12:19	11.5	10.78	39.9	11.5		3.0	4.0

RM	Unit	Habitat type	Sample date	Start time	Water temperature (C)	DO (ppm)	Specific conductivity (mS)	Horizontal visibility (ft)	Vertical visibility (ft)	Average depth (ft)	Maximum depth (ft)
42.3	126	Riffle	7-Mar	12:59	12.8	11.70	42.4	11.5		1.0	3.0
41.9	133	Run Head	3-Mar	16:49	10.0	10.25	39.9	8.0		2.5	4.0
41.8	134	Run Body	3-Mar	16:02	10.9	10.25	39.9	8.0		4.0	8.0
39.2	165	Pool Head	7-Mar	15:42	14.1	12.31	53.4	9.0		3.0	5.0
38.9	166	Pool Body	7-Mar	15:45	14.1	12.31	53.4	9.0	12.0	7.0	13.0
38.9	168	Riffle	8-Mar	11:00	12.1	10.65	48.9	8.5		1.5	3.5
38.8	172	Run Head	8-Mar	11:42	12.4	11.12	49.1	8.5		1.5	3.0
38.7	173	Run Body	8-Mar	11:28	12.4	11.12	49.1	8.5		2.0	3.0
38.5	179	Riffle	8-Mar	12:52	12.4	11.12	49.1	8.5		1.5	4.0

Table E-2. Water quality data for the sampling units selected for snorkel sampling, August 2010.

RM	Unit	Habitat type	Sample date	Start time	Water temperature (C)	DO (ppm)	Specific conductivity (mS)	Horizontal visibility (ft)	Vertical visibility (ft)	Average depth (ft)	Maximum depth (ft)
51.8	1	Pool Head	17-Aug	16:54	12.6	9.8	30.4	32.0	7.0	4.0	7.0
51.6	4	Pool Head	17-Aug	14:11	12.6	9.8	30.4	32.0	5.0	4.0	5.0
51.6	5	Pool Body	17-Aug	12:15	12.6	9.8	30.4	32.0	32.0	20.0	32.0
50.8	12	Run Body	18-Aug	15:28	13.1	11.0	29.1	31.5	8.0	6.0	8.0
50.6	14	Riffle	18-Aug	11:43	13.1	11.0	29.1	31.5	4.5	2.0	4.5
50.3	19	Run Head	18-Aug	10:58	12.7	11.2	28.8	27.3	9.0	5.0	9.0
49.9	24	Run Body	19-Aug	12:40	14.3	11.3	29.3	27.3	8.0	4.0	8.0
49.7	27	Pool Head	19-Aug	15:43	14.3	11.3	29.3	27.3	4.0	3.0	4.0
49.6	28	Pool Body	19-Aug	15:00	14.3	11.3	29.3	27.3	18.6	8.0	18.6
49.1	38	Run Head	20-Aug	14:15	14.2	11.2	29.7	25.0	2.5	2.0	2.5
48.4	45	Riffle	20-Aug	11:16	14.2	11.2	29.7	25.0	4.5	2.0	4.5
48.1	51	Run Body	20-Aug	15:25	16.4	13.1	29.4	25.0	8.0	6.0	8.0
48.0	53	Riffle	20-Aug	15:10	16.4	13.1	29.4	25.0	2.5	1.5	2.5
48.0	54	Pool Head	20-Aug	14:50	16.4	13.1	29.4	25.0	10.0	8.0	10.0
46.9	62	Run Head	21-Aug	12:30	13.9	11.8	30.4	20.5	4.5	3.0	4.5
45.3	81	Pool Body	21-Aug	14:40	15.3	12.7	31.1	20.5	19.5	10.0	19.5
45.1	83	Run Body	21-Aug	15:00	15.3	12.7	31.1	20.5	6.0	3.0	6.0
45.0	86	Pool Head	22-Aug	11:36	13.3	10.9	31.5	19.0	7.5	4.0	7.5
44.8	90	Run Head	22-Aug	12:16	13.3	10.9	31.5	19.0	2.0	0.5	2.0
44.5	101	Riffle	22-Aug	12:47	13.3	10.9	31.5	19.0	7.0	2.5	7.0
43.7	104	Pool Body	22-Aug	15:38	15.4	11.2	32.0	21.5	22.0	10.0	22.0
43.2	107	Riffle	22-Aug	17:00	15.4	11.2	32.0	21.5	6.0	1.5	6.0
42.7	123	Run Head	23-Aug	11:27	15.6	11.3	33.2	19.5	3.0	1.5	3.0
42.4	124	Run Body	23-Aug	11:38	15.6	11.3	33.2	19.5	4.5	3.0	4.5
40.3	150	Run Body	23-Aug	15:05	18.5	12.0	37.1	16.5	4.0	1.5	4.0
39.7	156	Riffle	23-Aug	16:18	18.5	12.0	37.1	16.5	2.0	1.0	2.0

RM	Unit	Habitat type	Sample date	Start time	Water temperature (C)	DO (ppm)	Specific conductivity (mS)	Horizontal visibility (ft)	Vertical visibility (ft)	Average depth (ft)	Maximum depth (ft)
39.6	157	Run Head	23-Aug	16:03	18.5	12.0	37.1	16.5	3.0	2.0	3.0
39.2	165	Pool Head	24-Aug	11:24	16.3	9.7	38.2	17.5	4.0	2.0	4.0
38.9	166	Pool Body	24-Aug	11:26	16.3	9.7	38.2	17.5	10.0	5.0	10.0
38.9	168	Riffle	24-Aug	10:57	16.3	9.7	38.2	17.5	3.5	1.5	3.5
38.8	171	Pool Body	24-Aug	10:23	16.3	9.7	38.2	17.5	13.0	9.0	13.0

Appendix F: Water Temperature Data




Figure F-1. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Riffle A7 (RM 50.8), February-March 2010.




Figure F-2. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Riffle 13B (RM 45.5), February-March 2010.




Figure F-3. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Roberts Ferry Bridge (RM 39.6), February-March 2010.




Figure F-4. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Ruddy Gravel (RM 36.5), February-March 2010.




Figure F-5. Average daily water temperature from thermographs, February-March 2010.




Figure F-6. Daily average, minimum, and maximum air temperature at the Modesto Airport, February-March 2010.




Figure F-7. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Riffle A7 (RM 50.8), July-August 2010.




Figure F-8. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Riffle 13B (RM 45.5), July-August 2010.




Figure F-9. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Roberts Ferry Bridge (RM 39.6), July-August 2010.




Figure F-10. Hourly, mean weekly average, and 7-day average of daily maximum temperatures at Ruddy Gravel (RM 36.5), July-August 2010.




Figure F-11. Average daily water temperature from thermographs, July-August 2010.




Figure F-12. Daily average, minimum, and maximum air temperature at the Modesto Airport, July-August 2010.

Appendix G: Fish Observation Data

Table G-1. *O. mykiss* observation data for the sampling units, March 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
51.6	4	Pool Head	M	1	1	0-50
51.6	4	Pool Head	M	2	0	--
51.6	4	Pool Head	M	3	1	0-50
51.6	4	Pool Head	M	3	1	400-450
51.6	5	Pool Body	M	1	1	400-450
51.6	5	Pool Body	M	1	2	550-600
51.6	5	Pool Body	M	2	2	400-450
51.6	5	Pool Body	M	3	1	400-450
50.9	11	Pool Body	S	1	0	--
50.8	12	Run Body	S	1	0	--
50.6	15	Run Head	M	1	1	350-400
50.6	15	Run Head	M	2	0	--
50.6	15	Run Head	M	3	0	--
50.5	16	Run Body	M	1	0	--
50.5	16	Run Body	M	2	0	--
50.5	16	Run Body	M	3	0	--
50.3	18	Riffle	S	1	0	--
50.3	19	Run Head	S	1	1	450-500
50.1	20	Run Body	S	1	0	--
50.1	22	Riffle	M	1	0	--
50.1	22	Riffle	M	2	0	--
50.1	22	Riffle	M	3	0	--
49.7	26	Riffle	M	1	1	250-300
49.7	26	Riffle	M	2	0	--
49.7	26	Riffle	M	3	2	250-300
49.7	27	Pool Head	S	1	0	--
49.6	28	Pool Body	M	1	1	400-450
49.6	28	Pool Body	M	2	0	--
49.6	28	Pool Body	M	3	1	400-450
48.8	42	Run Head	M	1	0	--
48.8	42	Run Head	M	2	0	--
48.8	42	Run Head	M	3	0	--
48.7	43	Run Body	S	1	0	--
48.0	54	Pool Head	S	1	0	--
45.9	70	Riffle	S	1	0	--
45.0	86	Pool Head	M	1	0	--
45.0	86	Pool Head	M	2	0	--
45.0	86	Pool Head	M	3	0	--
44.8	90	Run Head	S	1	0	--

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
44.7	93	Riffle	M	1	0	--
44.7	93	Riffle	M	2	0	--
44.7	93	Riffle	M	3	0	--
44.5	101	Riffle	S	1	0	--
43.7	104	Pool Body	S	1	0	--
43.0	111	Riffle	S	1	0	--
43.0	112	Pool Head	M	1	0	--
43.0	112	Pool Head	M	2	1	300-350
43.0	112	Pool Head	M	3	2	300-350
43.0	113	Pool Body	M	1	0	--
43.0	113	Pool Body	M	2	0	--
43.0	113	Pool Body	M	3	0	--
42.9	116	Run Body	M	1	0	--
42.9	116	Run Body	M	2	0	--
42.9	116	Run Body	M	3	0	--
42.9	119	Run Head	S	1	0	--
42.3	126	Riffle	S	1	1	350-400
41.9	133	Run Head	M	1	0	--
41.9	133	Run Head	M	2	0	--
41.9	133	Run Head	M	3	0	--
41.8	134	Run Body	S	1	0	--
39.2	165	Pool Head	S	1	0	--
38.9	166	Pool Body	S	1	0	--
38.9	168	Riffle	S	1	0	--
38.8	172	Run Head	S	1	0	--
38.7	173	Run Body	M	1	0	--
38.7	173	Run Body	M	2	0	--
38.7	173	Run Body	M	3	0	--
38.5	179	Riffle	S	1	1	400-450

Table G-2. *O. mykiss* observation data for the sampling units, August 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
51.8	1	Pool Head	M	1	10	250-300
51.8	1	Pool Head	M	1	6	300-350
51.8	1	Pool Head	M	1	1	400-450
51.8	1	Pool Head	M	2	1	100-150
51.8	1	Pool Head	M	2	4	200-250
51.8	1	Pool Head	M	2	6	250-300
51.8	1	Pool Head	M	2	3	300-350
51.8	1	Pool Head	M	2	2	350-400
51.8	1	Pool Head	M	3	7	200-250
51.8	1	Pool Head	M	3	9	250-300
51.8	1	Pool Head	M	3	1	300-350
51.6	4	Pool Head	M	1	4	250-300
51.6	4	Pool Head	M	1	1	300-350
51.6	4	Pool Head	M	1	2	350-400
51.6	4	Pool Head	M	1	2	400-450
51.6	4	Pool Head	M	1	1	450-500
51.6	4	Pool Head	M	2	2	250-300
51.6	4	Pool Head	M	2	3	300-350
51.6	4	Pool Head	M	2	1	350-400
51.6	4	Pool Head	M	2	1	400-450
51.6	4	Pool Head	M	3	2	300-350
51.6	5	Pool Body	M	1	1	200-250
51.6	5	Pool Body	M	1	2	200-250
51.6	5	Pool Body	M	1	2	200-250
51.6	5	Pool Body	M	1	1	250-300
51.6	5	Pool Body	M	1	4	300-350
51.6	5	Pool Body	M	2	2	100-150
51.6	5	Pool Body	M	2	1	150-200
51.6	5	Pool Body	M	2	1	200-250
51.6	5	Pool Body	M	2	2	250-300
51.6	5	Pool Body	M	2	1	300-350
51.6	5	Pool Body	M	2	1	350-400
51.6	5	Pool Body	M	2	1	400-450
51.6	5	Pool Body	M	2	1	400-450
51.6	5	Pool Body	M	3	2	100-150
51.6	5	Pool Body	M	3	2	150-200
51.6	5	Pool Body	M	3	1	300-350
51.6	5	Pool Body	M	3	1	350-400
51.6	5	Pool Body	M	3	1	400-450

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
51.6	5	Pool Body	M	3	1	400-450
51.6	5	Pool Body	M	3	1	450-500
50.8	12	Run Body	M	1	5	100-150
50.8	12	Run Body	M	1	1	100-150
50.8	12	Run Body	M	1	9	100-150
50.8	12	Run Body	M	1	1	150-200
50.8	12	Run Body	M	1	4	150-200
50.8	12	Run Body	M	1	2	200-250
50.8	12	Run Body	M	1	4	250-300
50.8	12	Run Body	M	1	14	300-350
50.8	12	Run Body	M	1	3	350-400
50.8	12	Run Body	M	1	4	350-400
50.8	12	Run Body	M	1	5	50-100
50.8	12	Run Body	M	2	10	100-150
50.8	12	Run Body	M	2	3	100-150
50.8	12	Run Body	M	2	1	100-150
50.8	12	Run Body	M	2	1	150-200
50.8	12	Run Body	M	2	10	150-200
50.8	12	Run Body	M	2	2	150-200
50.8	12	Run Body	M	2	1	200-250
50.8	12	Run Body	M	2	1	200-250
50.8	12	Run Body	M	2	10	250-300
50.8	12	Run Body	M	2	2	300-350
50.8	12	Run Body	M	2	7	300-350
50.8	12	Run Body	M	2	5	300-350
50.8	12	Run Body	M	2	10	300-350
50.8	12	Run Body	M	2	5	350-400
50.8	12	Run Body	M	2	5	350-400
50.8	12	Run Body	M	3	5	100-150
50.8	12	Run Body	M	3	5	100-150
50.8	12	Run Body	M	3	10	100-150
50.8	12	Run Body	M	3	3	100-150
50.8	12	Run Body	M	3	1	150-200
50.8	12	Run Body	M	3	2	150-200
50.8	12	Run Body	M	3	1	200-250
50.8	12	Run Body	M	3	2	250-300
50.8	12	Run Body	M	3	10	250-300
50.8	12	Run Body	M	3	4	300-350
50.8	12	Run Body	M	3	5	300-350
50.8	12	Run Body	M	3	3	350-400
50.8	12	Run Body	M	3	1	400-450

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
50.8	12	Run Body	M	3	5	50-100
50.8	12	Run Body	M	3	45	50-100
50.6	14	Riffle	M	1	25	100-150
50.6	14	Riffle	M	1	4	100-150
50.6	14	Riffle	M	1	6	150-200
50.6	14	Riffle	M	1	13	150-200
50.6	14	Riffle	M	1	4	200-250
50.6	14	Riffle	M	1	6	200-250
50.6	14	Riffle	M	1	3	250-300
50.6	14	Riffle	M	1	1	250-300
50.6	14	Riffle	M	1	1	300-350
50.6	14	Riffle	M	1	1	50-100
50.6	14	Riffle	M	2	6	100-150
50.6	14	Riffle	M	2	35	100-150
50.6	14	Riffle	M	2	4	100-150
50.6	14	Riffle	M	2	10	100-150
50.6	14	Riffle	M	2	5	100-150
50.6	14	Riffle	M	2	5	150-200
50.6	14	Riffle	M	2	4	150-200
50.6	14	Riffle	M	2	6	150-200
50.6	14	Riffle	M	2	2	200-250
50.6	14	Riffle	M	2	1	200-250
50.6	14	Riffle	M	2	4	200-250
50.6	14	Riffle	M	2	1	200-250
50.6	14	Riffle	M	2	3	300-350
50.6	14	Riffle	M	2	1	50-100
50.6	14	Riffle	M	2	4	50-100
50.6	14	Riffle	M	3	18	100-150
50.6	14	Riffle	M	3	21	100-150
50.6	14	Riffle	M	3	15	100-150
50.6	14	Riffle	M	3	3	150-200
50.6	14	Riffle	M	3	11	150-200
50.6	14	Riffle	M	3	9	150-200
50.6	14	Riffle	M	3	5	150-200
50.6	14	Riffle	M	3	3	200-250
50.6	14	Riffle	M	3	5	200-250
50.6	14	Riffle	M	3	2	250-300
50.6	14	Riffle	M	3	2	300-350
50.6	14	Riffle	M	3	2	350-400
50.6	14	Riffle	M	3	1	50-100
50.6	14	Riffle	M	3	2	50-100

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
50.6	14	Riffle	M	3	3	50-100
50.3	19	Run Head	M	1	5	100-150
50.3	19	Run Head	M	1	1	100-150
50.3	19	Run Head	M	1	3	150-200
50.3	19	Run Head	M	1	3	250-300
50.3	19	Run Head	M	1	2	300-350
50.3	19	Run Head	M	2	5	100-150
50.3	19	Run Head	M	2	5	150-200
50.3	19	Run Head	M	2	5	200-250
50.3	19	Run Head	M	2	7	300-350
50.3	19	Run Head	M	3	5	150-200
50.3	19	Run Head	M	3	3	250-300
50.3	19	Run Head	M	3	7	300-350
49.9	24	Run Body	S	1	3	100-150
49.9	24	Run Body	S	1	3	100-150
49.9	24	Run Body	S	1	1	100-150
49.9	24	Run Body	S	1	4	150-200
49.9	24	Run Body	S	1	1	200-250
49.9	24	Run Body	S	1	2	250-300
49.9	24	Run Body	S	1	11	300-350
49.9	24	Run Body	S	1	2	300-350
49.9	24	Run Body	S	1	4	350-400
49.7	27	Pool Head	M	1	3	100-150
49.7	27	Pool Head	M	1	4	150-200
49.7	27	Pool Head	M	1	4	150-200
49.7	27	Pool Head	M	1	1	200-250
49.7	27	Pool Head	M	1	1	200-250
49.7	27	Pool Head	M	1	1	250-300
49.7	27	Pool Head	M	2	3	100-150
49.7	27	Pool Head	M	2	4	100-150
49.7	27	Pool Head	M	2	4	150-200
49.7	27	Pool Head	M	2	5	150-200
49.7	27	Pool Head	M	2	3	150-200
49.7	27	Pool Head	M	2	1	200-250
49.7	27	Pool Head	M	2	1	200-250
49.7	27	Pool Head	M	2	1	300-350
49.7	27	Pool Head	M	2	2	50-100
49.7	27	Pool Head	M	3	4	100-150
49.7	27	Pool Head	M	3	5	150-200
49.7	27	Pool Head	M	3	3	150-200
49.7	27	Pool Head	M	3	3	150-200

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
49.7	27	Pool Head	M	3	1	200-250
49.7	27	Pool Head	M	3	1	200-250
49.7	27	Pool Head	M	3	3	50-100
49.6	28	Pool Body	M	1	1	100-150
49.6	28	Pool Body	M	1	3	250-300
49.6	28	Pool Body	M	1	3	300-350
49.6	28	Pool Body	M	1	1	300-350
49.6	28	Pool Body	M	1	1	350-400
49.6	28	Pool Body	M	1	1	350-400
49.6	28	Pool Body	M	2	2	100-150
49.6	28	Pool Body	M	2	4	150-200
49.6	28	Pool Body	M	2	2	200-250
49.6	28	Pool Body	M	2	3	250-300
49.6	28	Pool Body	M	2	5	250-300
49.6	28	Pool Body	M	2	5	300-350
49.6	28	Pool Body	M	2	2	350-400
49.6	28	Pool Body	M	2	1	350-400
49.6	28	Pool Body	M	3	2	100-150
49.6	28	Pool Body	M	3	2	150-200
49.6	28	Pool Body	M	3	3	250-300
49.6	28	Pool Body	M	3	1	250-300
49.6	28	Pool Body	M	3	5	300-350
49.1	38	Run Head	S	1	1	100-150
48.4	45	Riffle	S	1	11	100-150
48.4	45	Riffle	S	1	8	100-150
48.4	45	Riffle	S	1	7	100-150
48.4	45	Riffle	S	1	1	150-200
48.4	45	Riffle	S	1	4	150-200
48.4	45	Riffle	S	1	8	50-100
48.4	45	Riffle	S	1	1	50-100
48.1	51	Run Body	M	1	8	100-150
48.1	51	Run Body	M	1	8	100-150
48.1	51	Run Body	M	1	1	150-200
48.1	51	Run Body	M	1	3	150-200
48.1	51	Run Body	M	1	1	300-350
48.1	51	Run Body	M	1	1	350-400
48.1	51	Run Body	M	2	5	100-150
48.1	51	Run Body	M	2	10	100-150
48.1	51	Run Body	M	2	2	150-200
48.1	51	Run Body	M	2	2	150-200
48.1	51	Run Body	M	2	1	300-350

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
48.1	51	Run Body	M	2	1	350-400
48.1	51	Run Body	M	3	6	100-150
48.1	51	Run Body	M	3	1	150-200
48.1	51	Run Body	M	3	1	150-200
48.1	51	Run Body	M	3	1	200-250
48.1	51	Run Body	M	3	1	250-300
48.1	51	Run Body	M	3	1	300-350
48.1	51	Run Body	M	3	1	350-400
48.0	53	Riffle	S	1	2	100-150
48.0	53	Riffle	S	1	2	100-150
48.0	53	Riffle	S	1	1	350-400
48.0	54	Pool Head	S	1	6	100-150
48.0	54	Pool Head	S	1	4	150-200
48.0	54	Pool Head	S	1	1	150-200
48.0	54	Pool Head	S	1	1	200-250
48.0	54	Pool Head	S	1	1	300-350
48.0	54	Pool Head	S	1	2	300-350
46.9	62	Run Head	M	1	3	100-150
46.9	62	Run Head	M	1	5	150-200
46.9	62	Run Head	M	1	1	200-250
46.9	62	Run Head	M	1	2	300-350
46.9	62	Run Head	M	1	1	350-400
46.9	62	Run Head	M	2	1	100-150
46.9	62	Run Head	M	2	2	100-150
46.9	62	Run Head	M	2	5	150-200
46.9	62	Run Head	M	2	2	200-250
46.9	62	Run Head	M	2	1	200-250
46.9	62	Run Head	M	3	5	100-150
46.9	62	Run Head	M	3	8	150-200
46.9	62	Run Head	M	3	1	200-250
45.3	81	Pool Body	S	1	0	--
45.1	83	Run Body	S	1	12	100-150
45.1	83	Run Body	S	1	1	100-150
45.1	83	Run Body	S	1	1	150-200
45.1	83	Run Body	S	1	8	150-200
45.1	83	Run Body	S	1	3	200-250
45.1	83	Run Body	S	1	1	300-350
45.1	83	Run Body	S	1	1	300-350
45.1	83	Run Body	S	1	3	300-350
45.0	86	Pool Head	S	1	7	100-150
45.0	86	Pool Head	S	1	1	100-150

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
45.0	86	Pool Head	S	1	2	150-200
45.0	86	Pool Head	S	1	5	150-200
45.0	86	Pool Head	S	1	1	150-200
45.0	86	Pool Head	S	1	3	150-200
45.0	86	Pool Head	S	1	3	200-250
45.0	86	Pool Head	S	1	2	250-300
45.0	86	Pool Head	S	1	1	250-300
45.0	86	Pool Head	S	1	2	250-300
45.0	86	Pool Head	S	1	1	300-350
45.0	86	Pool Head	S	1	1	300-350
44.8	90	Run Head	S	1	0	--
44.5	101	Riffle	M	1	10	100-150
44.5	101	Riffle	M	1	5	100-150
44.5	101	Riffle	M	1	1	150-200
44.5	101	Riffle	M	1	5	150-200
44.5	101	Riffle	M	1	3	150-200
44.5	101	Riffle	M	1	1	200-250
44.5	101	Riffle	M	1	1	200-250
44.5	101	Riffle	M	1	1	250-300
44.5	101	Riffle	M	2	3	100-150
44.5	101	Riffle	M	2	2	100-150
44.5	101	Riffle	M	2	4	100-150
44.5	101	Riffle	M	2	2	150-200
44.5	101	Riffle	M	2	2	150-200
44.5	101	Riffle	M	2	9	150-200
44.5	101	Riffle	M	2	1	200-250
44.5	101	Riffle	M	2	1	200-250
44.5	101	Riffle	M	3	5	100-150
44.5	101	Riffle	M	3	1	100-150
44.5	101	Riffle	M	3	3	100-150
44.5	101	Riffle	M	3	2	150-200
44.5	101	Riffle	M	3	3	150-200
44.5	101	Riffle	M	3	6	150-200
44.5	101	Riffle	M	3	1	200-250
44.5	101	Riffle	M	3	1	200-250
44.5	101	Riffle	M	3	1	250-300
43.7	104	Pool Body	S	1	0	--
43.7	104	Pool Body	S	1	0	--
43.2	107	Riffle	M	1	8	100-150
43.2	107	Riffle	M	1	5	100-150
43.2	107	Riffle	M	1	3	100-150

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
43.2	107	Riffle	M	1	4	150-200
43.2	107	Riffle	M	1	1	150-200
43.2	107	Riffle	M	1	3	150-200
43.2	107	Riffle	M	1	1	200-250
43.2	107	Riffle	M	1	1	250-300
43.2	107	Riffle	M	1	1	300-350
43.2	107	Riffle	M	2	7	100-150
43.2	107	Riffle	M	2	4	100-150
43.2	107	Riffle	M	2	8	100-150
43.2	107	Riffle	M	2	1	150-200
43.2	107	Riffle	M	2	3	150-200
43.2	107	Riffle	M	2	3	150-200
43.2	107	Riffle	M	2	1	150-200
43.2	107	Riffle	M	2	2	200-250
43.2	107	Riffle	M	2	1	200-250
43.2	107	Riffle	M	2	1	250-300
43.2	107	Riffle	M	2	1	300-350
43.2	107	Riffle	M	2	1	300-350
43.2	107	Riffle	M	3	4	100-150
43.2	107	Riffle	M	3	1	100-150
43.2	107	Riffle	M	3	6	100-150
43.2	107	Riffle	M	3	3	100-150
43.2	107	Riffle	M	3	2	150-200
43.2	107	Riffle	M	3	1	150-200
43.2	107	Riffle	M	3	2	150-200
43.2	107	Riffle	M	3	1	150-200
43.2	107	Riffle	M	3	1	200-250
43.2	107	Riffle	M	3	1	200-250
43.2	107	Riffle	M	3	1	250-300
43.2	107	Riffle	M	3	1	300-350
42.7	123	Run Head	S	1	0	--
42.4	124	Run Body	M	1	11	100-150
42.4	124	Run Body	M	1	10	100-150
42.4	124	Run Body	M	1	2	150-200
42.4	124	Run Body	M	1	2	50-100
42.4	124	Run Body	M	2	9	100-150
42.4	124	Run Body	M	2	5	100-150
42.4	124	Run Body	M	2	2	150-200
42.4	124	Run Body	M	2	3	150-200
42.4	124	Run Body	M	2	7	50-100
42.4	124	Run Body	M	3	15	100-150

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
42.4	124	Run Body	M	3	4	100-150
42.4	124	Run Body	M	3	2	150-200
42.4	124	Run Body	M	3	3	150-200
42.4	124	Run Body	M	3	1	250-300
42.4	124	Run Body	M	3	2	50-100
40.3	150	Run Body	S	1	2	100-150
40.3	150	Run Body	S	1	2	150-200
40.3	150	Run Body	S	1	1	150-200
40.3	150	Run Body	S	1	1	200-250
39.7	156	Riffle	S	1	1	100-150
39.7	156	Riffle	S	1	1	150-200
39.6	157	Run Head	M	1	0	--
39.6	157	Run Head	M	2	0	--
39.6	157	Run Head	M	3	0	--
39.2	165	Pool Head	S	1	0	--
38.9	166	Pool Body	S	1	0	--
38.9	168	Riffle	S	1	0	--
38.8	171	Pool Body	M	1	0	--
38.8	171	Pool Body	M	2	0	--
38.8	171	Pool Body	M	3	0	--

Table G-3. *O. tshawytscha* observation data for the sampling units, March 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
51.6	4	Pool Head	M	1	2	0-50
51.6	4	Pool Head	M	2	18	0-50
51.6	4	Pool Head	M	3	10	0-50
51.6	5	Pool Body	M	1	1	0-50
51.6	5	Pool Body	M	1	75	0-50
51.6	5	Pool Body	M	2	63	0-50
51.6	5	Pool Body	M	3	64	0-50
51.6	5	Pool Body	M	3	1	0-50
50.9	11	Pool Body	S	1	0	--
50.8	12	Run Body	S	1	0	--
50.6	15	Run Head	M	1	0	--
50.6	15	Run Head	M	2	0	--
50.6	15	Run Head	M	3	0	--
50.5	16	Run Body	M	1	0	--
50.5	16	Run Body	M	2	0	--
50.5	16	Run Body	M	3	0	--
50.3	18	Riffle	S	1	135	0-50
50.3	18	Riffle	S	1	37	0-50
50.3	18	Riffle	S	1	7	50-100
50.3	18	Riffle	S	1	2	50-100
50.3	19	Run Head	S	1	0	--
50.1	20	Run Body	S	1	80	0-50
50.1	22	Riffle	M	1	8	0-50
50.1	22	Riffle	M	2	0	--
50.1	22	Riffle	M	3	0	--
49.7	26	Riffle	M	1	1	50-100
49.7	26	Riffle	M	2	0	--
49.7	26	Riffle	M	3	0	--
49.7	27	Pool Head	S	1	0	--
49.6	28	Pool Body	M	1	0	--
49.6	28	Pool Body	M	2	0	--
49.6	28	Pool Body	M	3	0	--
48.8	42	Run Head	M	1	0	--
48.8	42	Run Head	M	2	0	--
48.8	42	Run Head	M	3	0	--
48.7	43	Run Body	S	1	0	--
48.0	54	Pool Head	S	1	0	--
45.9	70	Riffle	S	1	40	0-50
45.9	70	Riffle	S	1	1	0-50

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
45.9	70	Riffle	S	1	25	50-100
45.0	86	Pool Head	M	1	0	--
45.0	86	Pool Head	M	2	0	--
45.0	86	Pool Head	M	3	0	--
44.8	90	Run Head	S	1	0	--
44.7	93	Riffle	M	1	2	0-50
44.7	93	Riffle	M	1	1	50-100
44.7	93	Riffle	M	2	3	0-50
44.7	93	Riffle	M	2	11	50-100
44.7	93	Riffle	M	3	6	0-50
44.7	93	Riffle	M	3	16	50-100
44.5	101	Riffle	S	1	1	0-50
43.7	104	Pool Body	S	1	0	--
43.0	111	Riffle	S	1	2	0-50
43.0	112	Pool Head	M	1	15	0-50
43.0	112	Pool Head	M	1	15	50-100
43.0	112	Pool Head	M	2	15	0-50
43.0	112	Pool Head	M	2	15	50-100
43.0	112	Pool Head	M	3	15	0-50
43.0	112	Pool Head	M	3	15	50-100
43.0	113	Pool Body	M	1	0	--
43.0	113	Pool Body	M	2	0	--
43.0	113	Pool Body	M	3	0	--
42.9	116	Run Body	M	1	20	0-50
42.9	116	Run Body	M	1	7	50-100
42.9	116	Run Body	M	1	37	50-100
42.9	116	Run Body	M	2	14	0-50
42.9	116	Run Body	M	2	6	50-100
42.9	116	Run Body	M	3	7	0-50
42.9	116	Run Body	M	3	16	0-50
42.9	116	Run Body	M	3	7	50-100
42.9	119	Run Head	S	1	0	--
42.3	126	Riffle	S	1	2	0-50
42.3	126	Riffle	S	1	10	50-100
41.9	133	Run Head	M	1	0	--
41.9	133	Run Head	M	2	0	--
41.9	133	Run Head	M	3	0	--
41.8	134	Run Body	S	1	1	0-50
39.2	165	Pool Head	S	1	0	--
38.9	166	Pool Body	S	1	0	--
38.9	168	Riffle	S	1	0	--

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
38.8	172	Run Head	S	1	8	0-50
38.8	172	Run Head	S	1	3	50-100
38.7	173	Run Body	M	1	1	0-50
38.7	173	Run Body	M	2	0	--
38.7	173	Run Body	M	3	0	--
38.5	179	Riffle	S	1	0	--

Table G-4. *O. tshawytscha* observation data for the sampling units, August 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
51.8	1	Pool Head	M	1	1	600-700
51.8	1	Pool Head	M	1	3	700-800
51.8	1	Pool Head	M	1	1	900-1000
51.8	1	Pool Head	M	2	2	600-700
51.8	1	Pool Head	M	2	3	700-800
51.8	1	Pool Head	M	2	1	900-1000
51.8	1	Pool Head	M	3	1	600-700
51.8	1	Pool Head	M	3	3	700-800
51.8	1	Pool Head	M	3	1	900-1000
51.6	4	Pool Head	M	1	0	--
51.6	4	Pool Head	M	2	0	--
51.6	4	Pool Head	M	3	0	--
51.6	5	Pool Body	M	1	87	50-100
51.6	5	Pool Body	M	2	76	50-100
51.6	5	Pool Body	M	3	72	50-100
50.8	12	Run Body	M	1	4	0-50
50.8	12	Run Body	M	1	133	0-50
50.8	12	Run Body	M	1	5	50-100
50.8	12	Run Body	M	1	2	50-100
50.8	12	Run Body	M	2	7	0-50
50.8	12	Run Body	M	2	112	0-50
50.8	12	Run Body	M	2	10	100-150
50.8	12	Run Body	M	2	23	50-100
50.8	12	Run Body	M	2	5	50-100
50.8	12	Run Body	M	2	1	50-100
50.8	12	Run Body	M	3	148	0-50
50.8	12	Run Body	M	3	4	100-150
50.8	12	Run Body	M	3	10	100-150
50.8	12	Run Body	M	3	5	50-100
50.8	12	Run Body	M	3	8	50-100
50.6	14	Riffle	M	1	62	0-50
50.6	14	Riffle	M	1	32	0-50
50.6	14	Riffle	M	1	1	100-150
50.6	14	Riffle	M	1	3	100-150
50.6	14	Riffle	M	1	11	50-100
50.6	14	Riffle	M	1	7	50-100
50.6	14	Riffle	M	1	4	50-100
50.6	14	Riffle	M	2	39	0-50
50.6	14	Riffle	M	2	60	0-50

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
50.6	14	Riffle	M	2	4	100-150
50.6	14	Riffle	M	2	5	50-100
50.6	14	Riffle	M	2	7	50-100
50.6	14	Riffle	M	2	4	50-100
50.6	14	Riffle	M	3	38	0-50
50.6	14	Riffle	M	3	72	0-50
50.6	14	Riffle	M	3	2	100-150
50.6	14	Riffle	M	3	3	50-100
50.6	14	Riffle	M	3	28	50-100
50.3	19	Run Head	M	1	7	0-50
50.3	19	Run Head	M	1	1	0-50
50.3	19	Run Head	M	1	10	100-150
50.3	19	Run Head	M	1	40	50-100
50.3	19	Run Head	M	1	1	600-650
50.3	19	Run Head	M	2	9	0-50
50.3	19	Run Head	M	2	20	100-150
50.3	19	Run Head	M	2	30	50-100
50.3	19	Run Head	M	3	8	0-50
50.3	19	Run Head	M	3	1	0-50
50.3	19	Run Head	M	3	20	100-150
50.3	19	Run Head	M	3	30	50-100
49.9	24	Run Body	S	1	50	0-50
49.9	24	Run Body	S	1	20	100-150
49.9	24	Run Body	S	1	12	100-150
49.9	24	Run Body	S	1	1	150-200
49.9	24	Run Body	S	1	30	50-100
49.9	24	Run Body	S	1	7	50-100
49.7	27	Pool Head	M	1	1	100-150
49.7	27	Pool Head	M	3	3	50-100
49.6	28	Pool Body	M	1	1	100-150
49.6	28	Pool Body	M	1	3	50-100
49.6	28	Pool Body	M	2	0	--
49.6	28	Pool Body	M	3	4	150-200
49.1	38	Run Head	S	1	0	--
48.4	45	Riffle	S	1	30	0-50
48.4	45	Riffle	S	1	19	100-150
48.4	45	Riffle	S	1	15	100-150
48.4	45	Riffle	S	1	18	100-150
48.4	45	Riffle	S	1	62	50-100
48.4	45	Riffle	S	1	42	50-100
48.1	51	Run Body	M	1	14	0-50

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
48.1	51	Run Body	M	1	4	100-150
48.1	51	Run Body	M	1	3	50-100
48.1	51	Run Body	M	2	8	0-50
48.1	51	Run Body	M	2	3	100-150
48.1	51	Run Body	M	2	3	50-100
48.1	51	Run Body	M	2	2	50-100
48.1	51	Run Body	M	2	17	50-100
48.1	51	Run Body	M	3	12	0-50
48.1	51	Run Body	M	3	2	100-150
48.1	51	Run Body	M	3	2	150-200
48.1	51	Run Body	M	3	18	50-100
48.0	53	Riffle	S	1	2	50-100
48.0	53	Riffle	S	1	2	50-100
48.0	54	Pool Head	S	1	2	50-100
46.9	62	Run Head	M	1	9	0-50
46.9	62	Run Head	M	1	2	100-150
46.9	62	Run Head	M	1	1	100-150
46.9	62	Run Head	M	1	5	50-100
46.9	62	Run Head	M	1	9	50-100
46.9	62	Run Head	M	2	10	0-50
46.9	62	Run Head	M	2	6	100-150
46.9	62	Run Head	M	2	3	100-150
46.9	62	Run Head	M	2	2	50-100
46.9	62	Run Head	M	2	10	50-100
46.9	62	Run Head	M	3	10	100-150
46.9	62	Run Head	M	3	17	50-100
46.9	62	Run Head	M	3	10	50-100
45.3	81	Pool Body	S	1	0	--
45.1	83	Run Body	S	1	8	100-150
45.1	83	Run Body	S	1	20	50-100
45.0	86	Pool Head	S	1	0	--
44.8	90	Run Head	S	1	1	50-100
44.5	101	Riffle	M	1	5	0-50
44.5	101	Riffle	M	1	1	100-150
44.5	101	Riffle	M	1	5	100-150
44.5	101	Riffle	M	1	4	50-100
44.5	101	Riffle	M	1	2	50-100
44.5	101	Riffle	M	1	25	50-100
44.5	101	Riffle	M	2	3	0-50
44.5	101	Riffle	M	2	8	100-150
44.5	101	Riffle	M	2	1	100-150

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
44.5	101	Riffle	M	2	2	100-150
44.5	101	Riffle	M	2	22	50-100
44.5	101	Riffle	M	2	1	50-100
44.5	101	Riffle	M	2	6	50-100
44.5	101	Riffle	M	3	4	0-50
44.5	101	Riffle	M	3	6	100-150
44.5	101	Riffle	M	3	1	100-150
44.5	101	Riffle	M	3	2	100-150
44.5	101	Riffle	M	3	7	50-100
44.5	101	Riffle	M	3	1	50-100
44.5	101	Riffle	M	3	23	50-100
43.2	107	Riffle	M	1	3	100-150
43.2	107	Riffle	M	1	14	50-100
43.2	107	Riffle	M	1	3	50-100
43.2	107	Riffle	M	1	1	50-100
43.2	107	Riffle	M	2	2	100-150
43.2	107	Riffle	M	2	3	50-100
43.2	107	Riffle	M	2	6	50-100
43.2	107	Riffle	M	3	1	100-150
43.2	107	Riffle	M	3	4	50-100
43.2	107	Riffle	M	3	3	50-100
43.2	107	Riffle	M	3	6	50-100
43.2	107	Riffle	M	3	1	50-100
42.7	123	Run Head	S	1	0	--
42.4	124	Run Body	M	1	10	100-150
42.4	124	Run Body	M	1	1	100-150
42.4	124	Run Body	M	1	1	50-100
42.4	124	Run Body	M	1	9	50-100
42.4	124	Run Body	M	2	1	100-150
42.4	124	Run Body	M	2	4	100-150
42.4	124	Run Body	M	2	4	50-100
42.4	124	Run Body	M	2	7	50-100
42.4	124	Run Body	M	3	5	100-150
42.4	124	Run Body	M	3	1	50-100
42.4	124	Run Body	M	3	18	50-100
40.3	150	Run Body	S	1	0	--
39.7	156	Riffle	S	1	0	--
39.6	157	Run Head	M	1	0	--
39.6	157	Run Head	M	2	0	--
39.6	157	Run Head	M	3	0	--
39.2	165	Pool Head	S	1	0	--

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Sum of count	Size range
38.9	166	Pool Body	S	1	1	100-150
38.9	168	Riffle	S	1	2	100-150
38.8	171	Pool Body	M	1	0	--
38.8	171	Pool Body	M	2	0	--
38.8	171	Pool Body	M	3	0	--

Table G-5. Non-salmonid fish observation data for the sampling units, March 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
50.8	12	Run Body	S	1	Sacramento sucker	1	300-350
50.6	15	Run Head	M	1	Sacramento sucker	3	500-550
50.5	16	Run Body	M	1	Sacramento sucker	28	400-450
50.5	16	Run Body	M	1	Sacramento sucker	25	450-500
50.5	16	Run Body	M	1	Sacramento sucker	5	500-550
50.5	16	Run Body	M	2	Sacramento sucker	28	400-450
50.5	16	Run Body	M	2	Sacramento sucker	16	450-500
50.5	16	Run Body	M	2	Sacramento sucker	1	525
50.5	16	Run Body	M	2	Sacramento sucker	1	545
50.5	16	Run Body	M	3	Sacramento sucker	35	400-450
50.5	16	Run Body	M	3	Sacramento sucker	14	450-500
50.5	16	Run Body	M	3	Sacramento sucker	5	500-550
50.5	16	Run Body	M	3	Sacramento sucker	1	525
50.3	18	Riffle	S	1	Sacramento sucker	6	300-350
50.3	18	Riffle	S	1	Sacramento sucker	4	350-400
50.3	18	Riffle	S	1	Sacramento sucker	10	400-450
50.3	18	Riffle	S	1	Sacramento sucker	5	450-500
50.1	20	Run Body	S	1	Sacramento sucker	3	300-350
50.1	20	Run Body	S	1	Sacramento sucker	6	350-400
50.1	20	Run Body	S	1	Sacramento sucker	10	400-450
50.1	20	Run Body	S	1	Sacramento sucker	8	450-500
50.1	22	Riffle	M	2	Sacramento sucker	1	425
49.7	26	Riffle	M	1	Sacramento sucker	2	300-350
49.7	26	Riffle	M	1	Sacramento sucker	1	350-400
49.7	26	Riffle	M	1	Sacramento sucker	4	400-450
49.7	26	Riffle	M	2	Sacramento sucker	3	400-450
49.7	26	Riffle	M	2	Sacramento sucker	3	450-500
49.7	26	Riffle	M	3	Sacramento sucker	1	450-500
49.7	27	Pool Head	S	1	Sacramento sucker	1	550-600
49.6	28	Pool Body	M	1	Sacramento sucker	7	450-500
49.6	28	Pool Body	M	1	Sacramento sucker	8	500-550
49.6	28	Pool Body	M	2	Sacramento sucker	4	450-500
49.6	28	Pool Body	M	2	Sacramento sucker	7	500-550
49.6	28	Pool Body	M	3	Sculpin sp.	1	75
49.6	28	Pool Body	M	3	Sacramento sucker	2	450-500
49.6	28	Pool Body	M	3	Sacramento sucker	5	500-550
48.8	42	Run Head	M	1	Sacramento sucker	6	300-350
48.7	43	Run Body	S	1	Sacramento sucker	8	300-350
48.7	43	Run Body	S	1	Sacramento sucker	8	450-500

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
48.0	54	Pool Head	S	1	Cyprinid sp.	10	0-50
48.0	54	Pool Head	S	1	Sacramento sucker	2	450-500
48.0	54	Pool Head	S	1	Sacramento sucker	1	740
45.9	70	Riffle	S	1	Catfish sp.	1	355
45.9	70	Riffle	S	1	Sacramento sucker	2	400-450
45.9	70	Riffle	S	1	Sacramento sucker	2	450-500
45.9	70	Riffle	S	1	Sacramento sucker	4	500-550
44.7	93	Riffle	M	2	Hardhead/Pikeminnow	5	0-50
44.7	93	Riffle	M	2	Hardhead/Pikeminnow	7	50-100
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	1	0-50
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	2	100-150
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	4	50-100
44.5	101	Riffle	S	1	Sacramento sucker	3	400-450
44.5	101	Riffle	S	1	Sacramento sucker	1	450-500
44.5	101	Riffle	S	1	Sacramento sucker	1	50-100
43.0	112	Pool Head	M	1	Sacramento sucker	1	305
43.0	112	Pool Head	M	1	Sacramento sucker	3	350-400
43.0	113	Pool Body	M	3	Largemouth bass	1	405
42.9	116	Run Body	M	1	Hardhead/Pikeminnow	1	125
42.9	116	Run Body	M	1	Sacramento sucker	1	450-500
42.9	116	Run Body	M	1	Sacramento sucker	1	475
42.9	116	Run Body	M	2	Sacramento sucker	2	400-450
42.9	116	Run Body	M	2	Sacramento sucker	3	450-500
42.9	116	Run Body	M	2	Sacramento sucker	1	500-550
42.9	116	Run Body	M	3	Hardhead/Pikeminnow	1	125
42.9	116	Run Body	M	3	Hardhead/Pikeminnow	2	150-200
42.3	126	Riffle	S	1	Sacramento sucker	3	450-500
41.9	133	Run Head	M	1	Sacramento sucker	4	450-500
41.9	133	Run Head	M	3	Sacramento sucker	1	500-550
41.8	134	Run Body	S	1	Sacramento sucker	14	400-450
41.8	134	Run Body	S	1	Sacramento sucker	15	450-500
41.8	134	Run Body	S	1	Sacramento sucker	19	500-550
41.8	134	Run Body	S	1	Sacramento sucker	1	650-700
38.9	166	Pool Body	S	1	Sacramento sucker	1	390
38.7	173	Run Body	M	1	Sculpin sp.	1	50-100
38.7	173	Run Body	M	1	Smallmouth bass	1	400-450
38.7	173	Run Body	M	2	Sacramento sucker	1	400-450
38.5	179	Riffle	S	1	Sacramento sucker	8	400-450
38.5	179	Riffle	S	1	Sacramento sucker	10	450-500
38.5	179	Riffle	S	1	Sacramento sucker	1	500-550

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
38.5	179	Riffle	S	1	Sacramento sucker	1	525
50.8	12	Run Body	S	1	Sacramento sucker	1	300-350
50.6	15	Run Head	M	1	Sacramento sucker	3	500-550
50.5	16	Run Body	M	1	Sacramento sucker	28	400-450
50.5	16	Run Body	M	1	Sacramento sucker	25	450-500
50.5	16	Run Body	M	1	Sacramento sucker	5	500-550
50.5	16	Run Body	M	2	Sacramento sucker	28	400-450
50.5	16	Run Body	M	2	Sacramento sucker	16	450-500
50.5	16	Run Body	M	2	Sacramento sucker	1	525
50.5	16	Run Body	M	2	Sacramento sucker	1	545
50.5	16	Run Body	M	3	Sacramento sucker	35	400-450
50.5	16	Run Body	M	3	Sacramento sucker	14	450-500
50.5	16	Run Body	M	3	Sacramento sucker	5	500-550
50.5	16	Run Body	M	3	Sacramento sucker	1	525
50.3	18	Riffle	S	1	Sacramento sucker	6	300-350
50.3	18	Riffle	S	1	Sacramento sucker	4	350-400
50.3	18	Riffle	S	1	Sacramento sucker	10	400-450
50.3	18	Riffle	S	1	Sacramento sucker	5	450-500
50.1	20	Run Body	S	1	Sacramento sucker	3	300-350
50.1	20	Run Body	S	1	Sacramento sucker	6	350-400
50.1	20	Run Body	S	1	Sacramento sucker	10	400-450
50.1	20	Run Body	S	1	Sacramento sucker	8	450-500
50.1	22	Riffle	M	2	Sacramento sucker	1	425
49.7	26	Riffle	M	1	Sacramento sucker	2	300-350
49.7	26	Riffle	M	1	Sacramento sucker	1	350-400
49.7	26	Riffle	M	1	Sacramento sucker	4	400-450
49.7	26	Riffle	M	2	Sacramento sucker	3	400-450
49.7	26	Riffle	M	2	Sacramento sucker	3	450-500
49.7	26	Riffle	M	3	Sacramento sucker	1	450-500
49.7	27	Pool Head	S	1	Sacramento sucker	1	550-600
49.6	28	Pool Body	M	1	Sacramento sucker	7	450-500
49.6	28	Pool Body	M	1	Sacramento sucker	8	500-550
49.6	28	Pool Body	M	2	Sacramento sucker	4	450-500
49.6	28	Pool Body	M	2	Sacramento sucker	7	500-550
49.6	28	Pool Body	M	3	Sculpin sp.	1	75
49.6	28	Pool Body	M	3	Sacramento sucker	2	450-500
49.6	28	Pool Body	M	3	Sacramento sucker	5	500-550
48.8	42	Run Head	M	1	Sacramento sucker	6	300-350
48.7	43	Run Body	S	1	Sacramento sucker	8	300-350
48.7	43	Run Body	S	1	Sacramento sucker	8	450-500

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
48.0	54	Pool Head	S	1	Cyprinid sp.	10	0-50
48.0	54	Pool Head	S	1	Sacramento sucker	2	450-500
48.0	54	Pool Head	S	1	Sacramento sucker	1	740
45.9	70	Riffle	S	1	Catfish sp.	1	355
45.9	70	Riffle	S	1	Sacramento sucker	2	400-450
45.9	70	Riffle	S	1	Sacramento sucker	2	450-500
45.9	70	Riffle	S	1	Sacramento sucker	4	500-550
44.7	93	Riffle	M	2	Hardhead/Pikeminnow	5	0-50
44.7	93	Riffle	M	2	Hardhead/Pikeminnow	7	50-100
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	1	0-50
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	2	100-150
44.7	93	Riffle	M	3	Hardhead/Pikeminnow	4	50-100
44.5	101	Riffle	S	1	Sacramento sucker	3	400-450
44.5	101	Riffle	S	1	Sacramento sucker	1	450-500
44.5	101	Riffle	S	1	Sacramento sucker	1	50-100
43.0	112	Pool Head	M	1	Sacramento sucker	1	305
43.0	112	Pool Head	M	1	Sacramento sucker	3	350-400
43.0	113	Pool Body	M	3	Largemouth bass	1	405
42.9	116	Run Body	M	1	Hardhead/Pikeminnow	1	125
42.9	116	Run Body	M	1	Sacramento sucker	1	450-500
42.9	116	Run Body	M	1	Sacramento sucker	1	475
42.9	116	Run Body	M	2	Sacramento sucker	2	400-450
42.9	116	Run Body	M	2	Sacramento sucker	3	450-500
42.9	116	Run Body	M	2	Sacramento sucker	1	500-550
42.9	116	Run Body	M	3	Hardhead/Pikeminnow	1	125
42.9	116	Run Body	M	3	Hardhead/Pikeminnow	2	150-200
42.3	126	Riffle	S	1	Sacramento sucker	3	450-500
41.9	133	Run Head	M	1	Sacramento sucker	4	450-500
41.9	133	Run Head	M	3	Sacramento sucker	1	500-550
41.8	134	Run Body	S	1	Sacramento sucker	14	400-450
41.8	134	Run Body	S	1	Sacramento sucker	15	450-500
41.8	134	Run Body	S	1	Sacramento sucker	19	500-550
41.8	134	Run Body	S	1	Sacramento sucker	1	650-700
38.9	166	Pool Body	S	1	Sacramento sucker	1	390
38.7	173	Run Body	M	1	Sculpin sp.	1	50-100
38.7	173	Run Body	M	1	Smallmouth bass	1	400-450
38.7	173	Run Body	M	2	Sacramento sucker	1	400-450
38.5	179	Riffle	S	1	Sacramento sucker	8	400-450
38.5	179	Riffle	S	1	Sacramento sucker	10	450-500
38.5	179	Riffle	S	1	Sacramento sucker	1	500-550

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
38.5	179	Riffle	S	1	Sacramento sucker	1	525

Table G-6. Non-salmonid fish observation data for the sampling units, August 2010.

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
51.8	1	Pool Head	Y	1	Hardhead/Pikeminnow	1	400-450
51.8	1	Pool Head	Y	1	Striped bass	1	300-350
51.6	5	Pool Body	Y	1	Striped bass	1	400-450
51.6	5	Pool Body	Y	2	Striped bass	2	450-500
50.8	12	Run Body	Y	1	Hardhead/Pikeminnow	1	0-50
50.8	12	Run Body	Y	1	Hardhead/Pikeminnow	1	300-350
50.8	12	Run Body	Y	1	Hardhead/Pikeminnow	5	400-450
50.8	12	Run Body	Y	1	Sculpin sp.	3	0-50
50.8	12	Run Body	Y	1	Sacramento sucker	2	300-350
50.8	12	Run Body	Y	1	Sacramento sucker	4	400-450
50.8	12	Run Body	Y	2	Hardhead/Pikeminnow	4	400-450
50.8	12	Run Body	Y	2	Sculpin sp.	3	0-50
50.8	12	Run Body	Y	2	Sacramento sucker	2	350-400
50.8	12	Run Body	Y	2	Sacramento sucker	3	400-450
50.8	12	Run Body	Y	3	Hardhead/Pikeminnow	3	400-450
50.8	12	Run Body	Y	3	Sculpin sp.	1	0-50
50.8	12	Run Body	Y	3	Sacramento sucker	64	0-50
50.8	12	Run Body	Y	3	Sacramento sucker	3	300-350
50.8	12	Run Body	Y	3	Sacramento sucker	4	400-450
50.6	14	Riffle	Y	1	Sculpin sp.	7	100-150
50.6	14	Riffle	Y	1	Sacramento sucker	6	300-350
50.6	14	Riffle	Y	2	Sculpin sp.	2	0-50
50.6	14	Riffle	Y	2	Sculpin sp.	2	100-150
50.6	14	Riffle	Y	2	Sculpin sp.	6	50-100
50.6	14	Riffle	Y	2	Sacramento sucker	2	0-50
50.6	14	Riffle	Y	3	Sculpin sp.	4	100-150
50.6	14	Riffle	Y	3	Sculpin sp.	2	50-100
50.6	14	Riffle	Y	3	Sacramento sucker	3	300-350
50.3	19	Run Head	Y	1	Lamprey sp.	1	150-200
50.3	19	Run Head	Y	1	Striped bass	1	400-450
50.3	19	Run Head	Y	1	Sacramento sucker	70	0-50
50.3	19	Run Head	Y	2	Striped bass	1	400-450
50.3	19	Run Head	Y	2	Sacramento sucker	65	0-50
50.3	19	Run Head	Y	3	Sacramento sucker	63	0-50
49.9	24	Run Body	N	1	Gambusia sp.	100	0-50
49.9	24	Run Body	N	1	Hardhead/Pikeminnow	40	0-50
49.9	24	Run Body	N	1	Hardhead/Pikeminnow	2	400-450
49.9	24	Run Body	N	1	Striped bass	1	300-350
49.9	24	Run Body	N	1	Sacramento sucker	35	0-50

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
49.9	24	Run Body	N	1	Sacramento sucker	15	300-350
49.9	24	Run Body	N	1	Sacramento sucker	17	350-400
49.9	24	Run Body	N	1	Sacramento sucker	100	400-500
49.9	24	Run Body	N	1	Sacramento sucker	6	50-100
49.7	27	Pool Head	Y	1	Sacramento sucker	1	350-400
49.7	27	Pool Head	Y	2	Sacramento sucker	1	350-400
49.6	28	Pool Body	Y	2	Sacramento sucker	7	0-50
49.6	28	Pool Body	Y	3	Sacramento sucker	10	0-50
49.1	38	Run Head	N	1	Gambusia sp.	3	0-50
49.1	38	Run Head	N	1	Sacramento sucker	40	0-50
48.4	45	Riffle	N	1	Gambusia sp.	3	0-50
48.4	45	Riffle	N	1	Sculpin sp.	2	100-150
48.4	45	Riffle	N	1	Sculpin sp.	1	50-100
48.1	51	Run Body	Y	1	Hardhead/Pikeminnow	8	0-50
48.1	51	Run Body	Y	1	Sacramento sucker	15	0-50
48.1	51	Run Body	Y	2	Sacramento sucker	10	0-50
48.1	51	Run Body	Y	3	Hardhead/Pikeminnow	1	200-250
48.1	51	Run Body	Y	3	Hardhead/Pikeminnow	1	350-400
48.1	51	Run Body	Y	3	Sacramento sucker	24	0-50
48.1	51	Run Body	Y	3	Sacramento sucker	1	300-350
48.0	53	Riffle	N	1	Sculpin sp.	1	100-150
48.0	53	Riffle	N	1	Sacramento sucker	1	350-400
48.0	53	Riffle	N	1	Sacramento sucker	3	50-100
48.0	54	Pool Head	N	1	Sacramento sucker	3	0-50
48.0	54	Pool Head	N	1	Sacramento sucker	1	300-350
46.9	62	Run Head	Y	1	Sacramento sucker	1	100-150
46.9	62	Run Head	Y	1	Sacramento sucker	4	50-100
46.9	62	Run Head	Y	2	Sacramento sucker	3	50-100
46.9	62	Run Head	Y	3	Sacramento sucker	1	100-150
46.9	62	Run Head	Y	3	Sacramento sucker	3	50-100
45.3	81	Pool Body	N	1	Hardhead/Pikeminnow	3	300-350
45.3	81	Pool Body	N	1	Hardhead/Pikeminnow	6	350-400
45.3	81	Pool Body	N	1	Hardhead/Pikeminnow	7	400-450
45.3	81	Pool Body	N	1	Hardhead/Pikeminnow	2	450-500
45.3	81	Pool Body	N	1	Hardhead/Pikeminnow	1	500-550
45.3	81	Pool Body	N	1	Sacramento sucker	16	300-350
45.3	81	Pool Body	N	1	Sacramento sucker	24	350-400
45.3	81	Pool Body	N	1	Sacramento sucker	13	400-450
45.3	81	Pool Body	N	1	Sacramento sucker	10	450-500
45.3	81	Pool Body	N	1	Sacramento sucker	10	500-550

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
45.1	83	Run Body	N	1	Hardhead/Pikeminnow	2	300-350
45.1	83	Run Body	N	1	Hardhead/Pikeminnow	3	350-400
45.1	83	Run Body	N	1	Sacramento sucker	1	0-50
45.1	83	Run Body	N	1	Sacramento sucker	21	300-350
45.1	83	Run Body	N	1	Sacramento sucker	77	350-400
45.1	83	Run Body	N	1	Sacramento sucker	16	400-450
45.0	86	Pool Head	N	1	Hardhead/Pikeminnow	9	150-200
45.0	86	Pool Head	N	1	Hardhead/Pikeminnow	8	200-250
45.0	86	Pool Head	N	1	Hardhead/Pikeminnow	15	250-300
45.0	86	Pool Head	N	1	Hardhead/Pikeminnow	3	300-350
45.0	86	Pool Head	N	1	Sacramento sucker	1	250-300
44.8	90	Run Head	N	1	Hardhead/Pikeminnow	1	50-100
44.5	101	Riffle	Y	1	Hardhead/Pikeminnow	13	150-200
44.5	101	Riffle	Y	1	Hardhead/Pikeminnow	9	200-250
44.5	101	Riffle	Y	1	Hardhead/Pikeminnow	3	300-350
44.5	101	Riffle	Y	1	Sacramento sucker	14	0-50
44.5	101	Riffle	Y	1	Sacramento sucker	1	100-150
44.5	101	Riffle	Y	1	Sacramento sucker	10	50-100
44.5	101	Riffle	Y	2	Hardhead/Pikeminnow	11	100-150
44.5	101	Riffle	Y	2	Hardhead/Pikeminnow	31	150-200
44.5	101	Riffle	Y	2	Hardhead/Pikeminnow	14	200-250
44.5	101	Riffle	Y	2	Hardhead/Pikeminnow	2	300-350
44.5	101	Riffle	Y	2	Sacramento sucker	12	0-50
44.5	101	Riffle	Y	2	Sacramento sucker	1	100-150
44.5	101	Riffle	Y	2	Sacramento sucker	3	200-250
44.5	101	Riffle	Y	2	Sacramento sucker	11	50-100
44.5	101	Riffle	Y	3	Hardhead/Pikeminnow	21	150-200
44.5	101	Riffle	Y	3	Hardhead/Pikeminnow	19	200-250
44.5	101	Riffle	Y	3	Hardhead/Pikeminnow	3	250-300
44.5	101	Riffle	Y	3	Hardhead/Pikeminnow	5	300-350
44.5	101	Riffle	Y	3	Sculpin sp.	1	0-50
44.5	101	Riffle	Y	3	Sacramento sucker	8	0-50
44.5	101	Riffle	Y	3	Sacramento sucker	3	200-250
44.5	101	Riffle	Y	3	Sacramento sucker	9	50-100
43.7	104	Pool Body	N	1	Largemouth bass	1	400-450
43.7	104	Pool Body	N	1	Hardhead/Pikeminnow	1	300-350
43.7	104	Pool Body	N	1	Striped bass	3	250-300
43.7	104	Pool Body	N	1	Striped bass	4	300-350
43.7	104	Pool Body	N	1	Striped bass	6	350-400
43.7	104	Pool Body	N	1	Striped bass	7	400-450

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
43.7	104	Pool Body	N	1	Striped bass	1	450-500
43.7	104	Pool Body	N	1	Sacramento sucker	25	300-350
43.7	104	Pool Body	N	1	Sacramento sucker	180	350-400
43.7	104	Pool Body	N	1	Sacramento sucker	110	400-450
43.7	104	Pool Body	N	1	Sacramento sucker	15	450-500
43.2	107	Riffle	Y	1	Hardhead/Pikeminnow	2	100-150
43.2	107	Riffle	Y	1	Hardhead/Pikeminnow	1	200-250
43.2	107	Riffle	Y	1	Sacramento sucker	6	0-50
43.2	107	Riffle	Y	2	Hardhead/Pikeminnow	4	100-150
43.2	107	Riffle	Y	2	Hardhead/Pikeminnow	6	150-200
43.2	107	Riffle	Y	2	Hardhead/Pikeminnow	3	200-250
43.2	107	Riffle	Y	2	Sacramento sucker	8	0-50
43.2	107	Riffle	Y	3	Hardhead/Pikeminnow	3	100-150
43.2	107	Riffle	Y	3	Hardhead/Pikeminnow	6	150-200
43.2	107	Riffle	Y	3	Hardhead/Pikeminnow	2	200-250
43.2	107	Riffle	Y	3	Hardhead/Pikeminnow	2	250-300
43.2	107	Riffle	Y	3	Sacramento sucker	3	0-50
42.4	124	Run Body	Y	1	Hardhead/Pikeminnow	41	150-200
42.4	124	Run Body	Y	1	Hardhead/Pikeminnow	3	200-250
42.4	124	Run Body	Y	1	Hardhead/Pikeminnow	3	250-300
42.4	124	Run Body	Y	1	Hardhead/Pikeminnow	1	300-350
42.4	124	Run Body	Y	1	Hardhead/Pikeminnow	4	450-500
42.4	124	Run Body	Y	1	Sacramento sucker	3	250-300
42.4	124	Run Body	Y	1	Sacramento sucker	38	300-350
42.4	124	Run Body	Y	1	Sacramento sucker	14	350-400
42.4	124	Run Body	Y	2	Hardhead/Pikeminnow	40	150-200
42.4	124	Run Body	Y	2	Hardhead/Pikeminnow	2	250-300
42.4	124	Run Body	Y	2	Hardhead/Pikeminnow	3	300-350
42.4	124	Run Body	Y	2	Hardhead/Pikeminnow	5	450-500
42.4	124	Run Body	Y	2	Smallmouth bass	1	250-300
42.4	124	Run Body	Y	2	Sacramento sucker	4	150-200
42.4	124	Run Body	Y	2	Sacramento sucker	6	200-250
42.4	124	Run Body	Y	2	Sacramento sucker	33	250-300
42.4	124	Run Body	Y	2	Sacramento sucker	124	300-350
42.4	124	Run Body	Y	2	Sacramento sucker	8	350-400
42.4	124	Run Body	Y	2	Sacramento sucker	5	400-450
42.4	124	Run Body	Y	3	Hardhead/Pikeminnow	2	150-200
42.4	124	Run Body	Y	3	Hardhead/Pikeminnow	3	300-350
42.4	124	Run Body	Y	3	Hardhead/Pikeminnow	1	350-400
42.4	124	Run Body	Y	3	Hardhead/Pikeminnow	3	450-500

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
42.4	124	Run Body	Y	3	Sacramento sucker	3	150-200
42.4	124	Run Body	Y	3	Sacramento sucker	5	250-300
42.4	124	Run Body	Y	3	Sacramento sucker	147	300-350
42.4	124	Run Body	Y	3	Sacramento sucker	12	350-400
42.4	124	Run Body	Y	3	Sacramento sucker	6	400-450
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	3	100-150
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	13	150-200
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	19	200-250
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	1	200-250
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	12	250-300
40.3	150	Run Body	N	1	Hardhead/Pikeminnow	1	250-300
39.7	156	Riffle	N	1	Hardhead/Pikeminnow	3	100-150
39.7	156	Riffle	N	1	Hardhead/Pikeminnow	3	150-200
39.7	156	Riffle	N	1	Sacramento sucker	150	0-50
39.7	156	Riffle	N	1	Sacramento sucker	1	400-450
39.7	156	Riffle	N	1	Sacramento sucker	15	50-100
39.6	157	Run Head	Y	1	Hardhead/Pikeminnow	1	150-200
39.6	157	Run Head	Y	1	Sacramento sucker	10	300-350
39.6	157	Run Head	Y	1	Sacramento sucker	10	350-400
39.6	157	Run Head	Y	1	Sacramento sucker	15	400-450
39.6	157	Run Head	Y	1	Sacramento sucker	40	450-500
39.6	157	Run Head	Y	2	Sacramento sucker	5	300-350
39.6	157	Run Head	Y	2	Sacramento sucker	10	350-400
39.6	157	Run Head	Y	2	Sacramento sucker	30	400-450
39.6	157	Run Head	Y	2	Sacramento sucker	30	450-500
39.6	157	Run Head	Y	3	Hardhead/Pikeminnow	2	250-300
39.6	157	Run Head	Y	3	Sacramento sucker	5	300-350
39.6	157	Run Head	Y	3	Sacramento sucker	10	350-400
39.6	157	Run Head	Y	3	Sacramento sucker	25	400-450
39.6	157	Run Head	Y	3	Sacramento sucker	21	450-500
38.9	166	Pool Body	N	1	Largemouth bass	1	400-450
38.9	166	Pool Body	N	1	Hardhead/Pikeminnow	1	150-200
38.9	166	Pool Body	N	1	Hardhead/Pikeminnow	15	200-250
38.9	166	Pool Body	N	1	Hardhead/Pikeminnow	3	250-300
38.9	166	Pool Body	N	1	Hardhead/Pikeminnow	4	350-400
38.9	166	Pool Body	N	1	Striped bass	1	450-500
38.9	166	Pool Body	N	1	Smallmouth bass	1	200-250
38.9	166	Pool Body	N	1	Sacramento sucker	2	300-350
38.9	166	Pool Body	N	1	Sacramento sucker	9	350-400
38.9	168	Riffle	N	1	Hardhead/Pikeminnow	1	250-300

RM	Unit	Habitat	Single (S) or multiple (M) pass	Pass	Species	Sum of count	Size range
38.8	171	Pool Body	Y	1	Sacramento sucker	1	200-250
